



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 974 359 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
26.01.2000 Bulletin 2000/04

(51) Int Cl.7: **A61K 38/24**, A61K 47/10,  
A61K 47/18

(21) Application number: **99305664.7**

(22) Date of filing: **16.07.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(72) Inventors:  
• **Hoffmann, James Arthur**  
**Greenwood, Indiana 46143 (US)**  
• **Lu, Jirong**  
**Indianapolis, Indiana 46326 (US)**

(30) Priority: **23.07.1998 US 93906 P**  
**30.07.1998 US 94611 P**  
**31.07.1998 US 94767 P**  
**01.09.1998 US 98711 P**  
**17.09.1998 US 100696 P**

(74) Representative: **Denholm, Anna Marie et al**  
**Eli Lilly and Company Limited,**  
**Lilly Research Center,**  
**Erl Wood Manor**  
**Windlesham, Surrey GU20 6PH (GB)**

(71) Applicant: **ELI LILLY AND COMPANY**  
**Indianapolis, Indiana 46285 (US)**

(54) **FSH and FSH variant formulations, products and methods**

(57) This invention relates to FSH or a FSH variant containing an alpha and beta subunit contained in formulations, and articles of manufacture. The invention provides advantageous new proteins and nucleic acids, multiuse pharmaceutical solutions, formulations and products of said proteins and nucleic acids where none

approved for commercial use had previously existed having such extended use indications. These products are particularly useful in therapeutic regimens for increasing serum levels of FSH or a FSH variant over a period of treatment. Thus, *inter alia*, the invention fills the need for convenient products of FSH or from a FSH variant.

**EP 0 974 359 A2**

## Description

[0001] This application claims the benefit of U.S. Provisionals 60/093906 filed July 23, 1998, 60/094611 filed July 30, 1998, 60/094767 filed July 31, 1998, 60/098711 filed September 1, 1998, 60/100696 filed September 17, 1998 each of which applications is entirely incorporated herein by reference.

[0002] This invention relates to new formulations, articles of manufacture and methods of using preparations of follicle stimulating hormone (FSH) or follicle stimulating hormone variants (FSH variants) known in the art. The invention also provides advantageous, multi-use and stable solutions and formulations and pharmaceutical products of which have not previously existed for therapeutic use. These formulations and products are particularly useful in therapeutic regimens for increasing serum levels of FSH or FSH variants over a period of treatment. Thus, *inter alia*, the invention fills the need for convenient stable and preserved solutions, formulations and products comprising FSH or FSH variants and using these formulations and products in the treatment of infertility.

[0003] FSH is indicated for use in infertility. The patients are administered daily or twice daily intramuscular ("IM") or subcutaneous ("SC") injections with dosage adjusted to response, usually ranging from 75-300 IU/day. The short half-life of FSH makes it necessary that the patients are given once or twice daily injections, extending to several days, depending on their ovarian or testicular response. A more stable formulation of FSH or of a FSH variant would provide improvements for use in therapy.

[0004] Although FSH has not been previously administered by approved modes of administration other than by IM or SC, other therapeutic proteins are expected to be administered over an extended number of days. Various delivery methods, including regular SC or IM injections over a period of time, transdermal patches, implants, osmotic pumps, micropumps, cartridges, pulmonary delivery systems, and the like, would be useful, e.g., in facilitating patient compliance, to reducing discomfort, or to facilitating administration. These extended treatment regimens generally require stable solutions or preservatives in the formulation.

[0005] Preservatives, in one aspect, prevent or minimize deleterious microbial contamination in the formulation. For conventional, non-protein therapeutics, antimicrobial preservative agents, such as chlorohexidine, phenol, benzyl alcohol, m-cresol, o-cresol, p-cresol, chlorocresol, phenylmercuric nitrate, thimerosal, benzoic acid, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal and various mixtures thereof, are often added to a liquid formulation to ensure sterility during shelf life and/or the multiple use regimen (Akers, MJ, Pharm. Technol. 8, 36-46, 1984; Gennaro, AR., Remington's Pharmaceutical Sciences, 17<sup>th</sup> edition., Mack, Easton, PA., 1278-1280, 1985). These preservatives as a class, however, tend to be detrimental to the stability of proteins. For example, a very effective preservative, m-cresol, has been reported to generally combine with and denature proteins (Development of Pharmaceutical Parenteral Dosage Forms, Bontempo, ed., Marcel Dekker, Inc., New York, New York, pp. 118-119, 1977). It also presents particular difficulty with the solution stability of hormones, such as human growth hormone (Maa YF and Hsu C, International Journal of Pharmaceutics, 140, pp. 155-168, 1996).

[0006] FSH is a member of the heterodimer, glycoprotein hormone family that includes thyroid stimulating hormone (TSH), chorionic gonadotropin (CG), and luteinizing hormone (LH) (Pierce JG and Parsons TF, Annu. Rev. Biochem., 50, 465-495, 1981; Baenziger and Green, Biochem. Biophys. Acta., 947, 287-306, 1988). The members of this family are heterodimers, held together generally by noncovalent interactions between the two different subunits. The human FSH (hFSH) heterodimer consists of (i) a mature 92 amino acid alpha subunit, which also is common to the other human family members (i.e., chorionic gonadotropin ("CG"), luteinizing hormone ("LH") and thyroid stimulating hormone ("TSH")); and (ii) a mature 111 amino acid beta subunit that is unique to FSH (Shome et al., J. Clin. Endocrinol. Metab. 39:187-205 (1974); Shome, et al., J. Prot. Chem, 7:325-339, 1988). The alpha and beta subunits bind non-covalently and, thus, the binding was thought to be more susceptible to protein destabilizing agents.

[0007] The native human and other mammalian FSH alpha and beta amino acid sequences and certain variants of these sequences were well known in the art prior to 1982 and cloning and expression of active human and other mammalian FSH in mammalian cells had been accomplished prior to 1985. The common gonadotropin alpha (or FSH alpha) subunit was sequenced from purified protein (Bellisario et al., J. Biol. Chem. 248:6796 (1973); Morgan et al., J. Biol. Chem. 250:5247 (1975)) and later cloned and expressed (Fiddes et al., Nature 281:351 (1979); Nature 286:684 (1981); J. Molec. Appl. Genet. 1:3-18 (1981)). The FSH beta subunit was sequenced from purified protein (Shome et al., J. Clin Endocrinol. Metab. 39:187 (1974); Saxena et al., J. Biol. Chem. 251:993 (1976)); (Sairam et al., Biochem. J. 197:541 (1981); Fujiki et al., Biochem Biophys. Acta 624:428 (1980)). Integrated Genetics reported the recombinant expression of a human CG (Biotechnology Newswatch (p. 3, June 20, 1983); Chemical and Engineering News 61:41 (Nov. 21, 1983); Genetic Technology News 3:9 (Dec. 12, 1983)) and in active form (Biotechnology Newswatch, Jan. 16, 1984)), and they also reported the successful cloning of FSH (Genetic Engineering Newsletter 4:4 (August 10, 1984)) and recombinant FSH produced in mammalian cells in active form (DNA 4:76 (published Jan. 16, 1985)). Amgen also reported the expression of an active bovine LH in CHO cells (Proc. Natl. Acad. Sci. USA 82:7280 (Nov. 1985)).

[0008] There is substantial evidence in the literature indicating that heterodimeric protein hormones can dissociate

under physiological or acidic conditions (Ryan, R.J., et al., Recent Progr. Hormone Res. 26:105-137; 1970; Strickland, TW and Puett, D, J. Biol. Chem., 257:2954-2960; 1982; Reichert LE and Ramsey RB, J. Biol. Chem., 250:3034-3040; 1975). Intact dimers are essential for biological activity and vital to secretion of FSH (Baenziger JU and Green ED, Biochem. Biophys. Acta, 947:287-306, 1988; Corless, et al., J. Cell Biol., 104:1173-1181, 1987). Attempts to counteract the instability of FSH include those where a single chain molecule is produced, incorporating two subunits into one stable molecule, and those where additional disulfides bonds are created to stabilize the interaction between the two subunits (Sugihara T., et al., J. Biol. Chem., 271:10445-10448, 1996; Heikoop J.C., et al., Nature Biotech, 15:658-62, 1997).

**[0009]** Donaldson, U.S. Patent No. 5,162,306, is directed to veterinary compositions comprising FSH and LH. These compositions are shown to be stable in thymol (5-methyl-2(1-methylethyl)phenol). Donaldson reports that thymol is one preservative in the list of preservatives in the U.S.P. XXI that will not damage glycoprotein hormones (U.S. Patent No. 5,162,306) in the disclosed SUPER-OV formulation.

**[0010]** Urinary derived FSH from postmenopausal women (hMG, marketed as Menotropin or Humagon™ by Organon and as urofollitropin or Metrodin™ by Serono) has been used as an injectable for over 30 years for the development of multiple follicles in ovulatory patients participating in Assisted Reproductive Technology (ART) programs and for the induction of ovulation in anovulatory infertile patients. (Fauser BCJM and Van Heusden AM, Endocrine Rev., 18, 71-106, 1997). More recently, CHO cell-derived recombinant human FSH (rhFSH) has become available (Keene J.L., et al., J. Biol. Chem., 264:4769-4752, 1989; Loumaye E., et al., Human Reprod. Update, 1:188-1999, 1995; Olijve W., et al., Mol. Hum. Reprod., 2:361-369, 1996).

**[0011]** Therapeutic FSH (either hMG or rhFSH) is currently supplied in a lyophilized form in ampules of 75 IU/vial and 150 IU/vial with a shelf life of one and a half to two years when stored at 2-25°C. Daily injections with starting doses of 75 IU or 150 IU are recommended for up to ten days to reach steady state concentrations of hFSH that are 1.5-2.5 times higher than that after a single dose administration. This dosing regime yields concentrations necessary for therapeutic efficacy, as FSH acts through a threshold mechanism (Schoemaker J., et al., Ann. NY. Acad. Sci. 687: 296-299, 1993). Depending on the patient's response, up to three cycles of treatment with increasing doses of FSH can be used. The patient or the partner is required to reconstitute a new vial of lyophilized material with diluent and administer it immediately after reconstitution (Package insert N1700101A, published in February 1996, for Fertinex™ (urofollitropin for injection, purified) for subcutaneous injection, by Serono Laboratories, Inc., Randolph, MA) on a daily basis. Any unused material is discarded.

**[0012]** Accordingly, there remains a need in the art to increase patient compliance via the development of stable formulations and preserved formulations of FSH or FSH variant proteins, and related articles of manufacture. These stable preparations are especially needed where extended treatments are required or advised, such as fertility treatments with FSH. There is also need to provide an FSH or FSH variant products that can be used and approved for multi-use administration over a period of twenty-four hours or greater. The invention also provides new stable solutions and formulations and preserved solutions and formulations of FSH and FSH variants and the related articles of manufacture that can also be used and approved for use over a period of twenty-four hours or greater.

**[0013]** This invention provides new formulations of FSH or FSH variants, their preparation, and their pharmaceutical or veterinary use in the treatment of fertility disorders and related articles of manufacture.

**[0014]** In one aspect, the invention provides preserved solutions and formulations comprising FSH or a FSH variant and a preservative selected from the group consisting of at least one phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, and derivatives or mixtures thereof in an aqueous diluent. Optionally, the preserved solutions and formulations contain a selected buffer and a salt.

**[0015]** In another aspect, the invention provides stable solutions and formulations comprising a FSH or a FSH variant and a selected buffer, which is preferably a phosphate buffer with saline or a chosen salt.

**[0016]** In another aspect, the invention provides for the treatment of infertility which comprises administering to a patient in need thereof the preserved formulation of FSH or a FSH variant in solution containing at least one preservative selected from the group consisting of a phenol, an m-cresol, a p-cresol, an o-cresol, a chlorocresol, a benzyl alcohol, an alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, at least one derivative thereof, or mixtures thereof.

**[0017]** In another aspect, the invention provides for the treatment of infertility which comprises administering to a patient in need thereof the stable formulation of FSH or a FSH variant in a stable solution, which is preferably a phosphate buffer with saline or a chosen salt.

**[0018]** Another aspect of the invention provides a process for preparing at least one multi-dose formulation of FSH or a FSH variant, comprising admixing FSH and at least one preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, derivatives thereof, or mixtures thereof in an aqueous diluent.

**[0019]** Another aspect of the invention provides a process for preparing at least one stable formulation of FSH or a FSH variant, comprising admixing FSH or a FSH variant in stable solution or formulation, which is preferable a phosphate buffer with saline or a chosen salt.

**[0020]** This invention also provides an article of manufacture for human pharmaceutical use, comprising packaging material and a vial comprising a solution of FSH or a FSH variant and a preservative, wherein said packaging material comprises a label which indicates that such solution may be held over a period of twenty-four hours or greater for use. The invention further comprises an article of manufacture for human pharmaceutical use, comprising packaging material, a first vial comprising lyophilized FSH or a FSH variant, and a second vial comprising a preservative, wherein said packaging material comprises a label which instructs a patient to reconstitute the FSH or a FSH variant in the preservative solution to form a solution which may be held over a period of twenty-four hours or greater for use under conditions as further described herein.

**[0021]** This invention also provides an article of manufacture for human pharmaceutical use, comprising packaging material and a vial comprising a solution of FSH or a FSH variant and stable solution or formulation, which is preferable a phosphate buffer with saline or a chosen salt, wherein said packaging material comprises a label which indicates that such solution may be held over a period of twenty-four hours or greater for use.

**[0022]** The invention further comprises an article of manufacture for human pharmaceutical use, comprising packaging material, a first vial comprising lyophilized FSH or a FSH variant, and a second vial comprising a preservative, wherein said packaging material comprises a label which instructs a patient to reconstitute the FSH or a FSH variant in the preservative solution to form a solution which may be held over a period of twenty-four hours or greater for use under conditions as further described herein.

**[0023]** This invention also provides an article of manufacture for human pharmaceutical use, comprising packaging material and a vial comprising a lyophilized FSH or a FSH variant and a second stable solution or formulation, which is preferable a phosphate buffer with saline or a chosen salt, wherein said packaging material comprises a label which instructs a patient to reconstitute the FSH or a FSH variant in the stable solution to form a solution which may be held over a period of twenty-four hours or greater for use under conditions as further described herein.

**[0024]** The present invention, in one aspect, provides recombinant and/or purified or isolated FSH or a FSH variant solutions and formulations, articles of manufacture and methods of use or treatment, and pharmaceutical products that are unexpectedly stable and/or are suitable for extended or multiple use.

**[0025]** These FSH or FSH variant solutions and formulations, articles of manufacture, methods of use and treatment using a FSH or a FSH variant, with improved or more suitable properties or stability, are useful for infertility treatment in women and/or men. These formulations, articles of manufacture, are additionally suitable for use in injectable and alternative delivery systems, e.g., but not limited to, nasal, pulmonary, transmucosal, transdermal, oral, subcutaneous, intramuscular or parenteral sustained release, dry, or liquid formulation. The FSH or a FSH variant solutions and formulations provided may also have increase *in vivo* potency compared to known commercial products, alone or in combination with at least one additional gonadotropin, by preventing or reducing loss of activity or stability, or by improving any aspect of the effectiveness or desirability of administration, e.g., by at least one of mode, frequency, dosage, comfort, ease of use, biological activity *in vitro* or *in vivo*, and the like.

**[0026]** All publications or patents cited herein are each entirely incorporated herein by reference as they show the state of the art at the time of the present invention or the filing dates of the related patent applications cited herein.

The following citations are entirely incorporated by reference: Ausubel, et al., ed., Current Protocols in Molecular Biology, John Wiley and Sons, NY (1987-1999); Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> Edition, Cold Spring Harbor, NY (1989); Harlow and Lane, Antibodies, a Laboratory Manual, Cold Spring Harbor, NY (1989); Colligan, et al., eds., Current Protocols in Immunology, John Wiley and Sons, N.Y. (1994-1999); Colligan et al., eds., Current Protocols in Protein Science, John Wiley and Sons, N.Y. (1998-1999).

**[0027]** Follicle stimulating hormone "FSH", whether produced recombinantly or isolated, and follicle stimulating hormone variants "FSH variants" as defined herein are well-known in the art. FSH as used herein refers to the FSH produced as a full length mature protein which includes, but are not limited to human FSH or "hFSH", whether produced recombinantly or isolated from human sources (see Shome B., et al., J. Prot. Chem., 7:325-339, 1988; Saxena B.B. and Rathnam P., J. Biol. Chem., 251:993-1005, 1976; Watkins, et al., DNA, 6:205-212, 1987; Shome B. and Parlow A.F., J. Clin. Endocrinol. Metab., 39(1):203-205, 1974; and Beck, et al., DNA, 4:76, 1985; U.S. 5,405,945, and U.S. 5,639,640—each citation incorporated by reference. The protein sequence of the human FSH alpha subunit is provided in SEQ ID NO: 5, and the protein sequence of the human FSH beta subunit is given in SEQ ID NO:6. Furthermore, various FSH variants are known or are understood from the art (see Shome, J. Clin. Endocrin. Metab 39:187 (1974); Saxena, J. Biol Chem 251(4):993-1005 (1976); 1978; Sairam et al., Biochem J 197:541 (1981); additionally see Closset Eur. J. Biochem. 86:115-120; Fujiki, J. Biol. Chem. 253:5363-5368 (1978); Sairam, Biochem. J. 197:541-552 (1981)—each citation independently incorporated by reference). \*\*\*Prior-art FSH beta subunits would include the Saxena sequence as well as a genus of sequences implicated in Sairam's discussion of (a) evolutionarily conserved amino acids and (b) well-known and characterized errors in sequencing. Further, those of skill in the art recognize that the substi-

tution of a prior art identified amino acid with (i) a chemically similar amino acid or (ii) an evolutionarily conserved amino acid would have no appreciable effect on the biological activity of an FSH heterodimer comprised of an hFSH beta subunit, thus modified.

**[0028]** In particular, Sairam's commentary on the Saxena hFSH sequence, as well as his discussion of amino acid substitutions identified between functional FSH molecules, defines a genus of FSH beta chain sequences in the prior art. More specifically, the 1981 Sairam publication identifies conserved amino acid sequences referring to publications by Saxena et al., Shome et al., Closset et al., and Fujiki et al. Sairam, *Biochem J* 197:541, 551 (1981). The prior art (1) evidences a preference for the FSH beta-chain sequence of Saxena over that of Shome; (2) addresses the issue of carboxy-terminal heterogeneity; (3) states that portions of the molecule affected by interspecies differences that are not essential for activity of the hormone and (4) highlights the guidance drawn from homologies between species and between the beta chains of the three, human glycoprotein hormones, FSH, LH and TSH.

**[0029]** C-terminal heterogeneity is reported for all the published sequences except for that of the porcine FSH-s, in which glutamic acid was the only C-terminal residue. For position 27, Saxena assigned one tryptophan residue to this position also found support in the evolutionary conservation demonstrated for a tryptophan at position 24 for FSH-B, among all prior art species. For positions 44 and 46, Saxena shows that, at position 44, the residue should be arginine instead of lysine and, at position 46, lysine instead of arginine. The porcine, equine and ovine sequences also reflected an evolutionary pressure to conserve the arginine at the position 44. The variations at three positions, 21, 22 and 44 involve a structurally conservative or evolutionarily-conserved ("homologous") substitutions, each of which possess bio-activity.

**[0030]** Each of the Sairam, Shome, and Closset references disclose residues isoleucine, serine at positions 21-22, while Saxena discloses leucine, threonine and Fujiki discloses isoleucine, threonine at these positions. Each of these disclosures is not only an evolutionarily conservative substitution, but also a structurally conservative substitution. The variation at position 41 between the aspartic acid disclosed by each of Sairam, Shome, Closset, and Fujiki and the asparagine disclosed by Saxena, Closset and Sairam involves two evolutionarily conserved residues, each of which provide bio-activity. These disclosures of conservative substitutions and evolutionarily conserved substitutions guide the skilled artisan to distinct FSH beta chain variants, within the hFSH-B chain genus.

**[0031]** The FSH variants referred to herein are the carboxy terminal deletions of the beta subunit that are shorter than the full length mature protein of SEQ ID NO:6. Carboxy terminal deletions of the human beta subunit are provided in SEQ IDS NOS: 11, 12, and 13. It is understood that the carboxy terminal variants of the beta chain form dimers with a known alpha subunit to form a FSH variant heterodimer. Additionally, a number of species of FSH are known, including but limited to porcine FSH (SEQ ID NOS: 7 and 8), horse FSH (SEQ ID NOS: 3 and 4), bovine FSH (SEQ ID NOS: 1 and 2), sheep FSH (SEQ ID NOS: 9 and 10), and those cited in Combarous Y., *Endocrine Reviews*, 13(4), 670-691, 1992; — herein incorporated by reference. Therein it is understood that one skilled in the art would be able to make and prepare other carboxy terminal variants from the given species as further provided herein.

**[0032]** FSH heterodimers or FSH variant heterodimers can be produced by any suitable method, such as recombinantly, by isolation or purification from natural sources as may be the case, or by chemical synthesis, or any combination thereof. Non-limiting examples FSH heterodimers and FSH variant heterodimers comprising one alpha subunit and one beta subunit include but are not limited to:

(a) : $\alpha$ -subunit: (SEQ ID NO:1)

FPDGEFTMQGCPCKLKENKYFSKPDAPYQCMGCCFSRAYPTPARSKKTMVLVPMK

ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:2)

RSCELTNITITVEKEECGFCISINTTWCAGYCYTRDLVYRDPARPNIQKTCTFKEL

VYETVKVPGCAHHADSLYTPVATECHCSKCDSDSTDCTVRGLGPSYCSFREIKE

(b): $\alpha$ -subunit:(SEQ ID NO:3)

FPDGEFTTQDCPECKLRENKYFFKLGVPPIYQCKGCCFSRAYPTPARSRKTM LVPKN

ITSESTCCVAKAFIRVTVMGNIKENHTQCYCSTCYHHKI

$\beta$ -subunit:(SEQ ID NO:4)

NSCELTNITIAVEKEGCGFCITINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL

VYETVKVPGCAHHADSLYTPVATACHCGKCNSDSTDCTVRGLGPSYCSFGDMKE

(c): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE

STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHHKS

$\beta$ -subunit:(SEQ ID NO:6)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL

VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMKE

(d): $\alpha$ -subunit:(SEQ ID NO:7)

FPDGEFTMQGCPECKLKENKYFSKLGAPIYQCMGCCFSRAYPTPARSKKTMLVPKN

ITSEATCCVAKAFTKATVMGNARVENHTECHCSTCYHHKS

$\beta$ -subunit:(SEQ ID NO:8)

NSCELTNITITVEKEECNFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL

VYETVKVPGCAHHADSLYTPVATECHCGKCDSDSTDCTVRGLGPSYCSFSEMKE

(e): $\alpha$ -subunit:(SEQ ID NO:9)

FPDGEFTMQGCPECKLKENKYFSKPDAPYQCMGCCFSRAYPTPARSKKTMLVPKN

ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHHKS

$\beta$ -subunit:(SEQ ID NO:10)

RSCELTNITITVEKEECRFCISINTTWCAGYCYTRDLVYKDPARPNIQKACTFKEL

VYETVKVPGCAHHADSLYTPVATECHCGKCDRDSTDCTVRGLGPSYCSFSDIRE

(f): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE

STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHHKS

$\beta$ -subunit:(SEQ ID NO:11)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL

VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGE

(g):  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:12)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEM

(h):  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:13)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMK

**[0033]** The use of the term "recombinant" refers to recombinant preparations of FSH or FSH variants through the use of recombinant DNA technology (e.g. Boine et al., Seminars in Reproductive Endocrinology 10, 45-50, 1992, and as generally further provided and exemplified herein). The sequences for genomic and cDNA clones are known for the alpha and beta subunits of several species (Fiddes, J.C., et al., J of Mol. and Applied Genetics, 1:3-18(1981); Esch F.S., et al. DNA 5:363-369(1986); Watkins P.C., et al., DNA 6:205-212(1987); Hirai T., et al., J. Mol. Endocrinol. 5: 147-158(1990); Maurer, R.A., et al., Mol. Endocrinol. 1:717-723(1987); Guzman K., et al., DNA Cell Biol. 10:593-601 (1991); Kumar TR, et al., Gene. 1995 Dec 12;166(2):335-6; Kumar TR, et al., Gene. 1995 Dec 12;166(2):333-4 - herein, each citation incorporated by reference). Several of the DNA sequences for alpha and beta subunits are provided as SEQ IDS: 14-20. Moreover, transfection of eucaryotic cells with the DNA sequences encoding a alpha and beta subunit, whether provided on one vector or on two vectors with each subunit having a separate promoter are capable of providing intact dimers, or by other methods understood in the art.

**[0034]** The FSH or a FSH variant used in accordance with the present invention may be produced not only by recombinant means, including from mammalian cell or transgenic preparations, but also may be purified from other biological sources, such as from urinary sources. Acceptable methodologies include those described in Hakola, K. Molecular and Cellular Endocrinology, 127:59-69, 1997; Keene, et al., J. Biol. Chem., 264:4769-4775, 1989; Cerpa-Poljak, et al., Endocrinology, 132:351-356, 1993; Dias, et al., J. Biol. Chem., 269:25289-25294, 1994; Flack, et al., J. Biol. Chem., 269:14015-14020, 1994; and Valove, et al., Endocrinology, 135:2657-2661, 1994, and U.S. Patent 3,119,740, herein entirely incorporated by reference.

**[0035]** "Substantially pure," used in reference to a peptide or protein, means separation from other cellular and non-cellular molecules, including other protein molecules. A substantially pure preparation would be about at least 85% pure; preferably about at least 95% pure. A "substantially pure" protein can be prepared by a variety of techniques, well known to the skilled artisan, including, for example, high pressure liquid chromatography (HPLC) and as further understood in the art or demonstrated herein.

**[0036]** The term "administer" or "administering" means to introduce a formulation of the present invention into the body of a patient in need thereof to treat a disease or condition.

**[0037]** The term "patient" means a mammal that is treated for a disease or condition. Patients are of, but not limited to, the following origin, human, ovine, porcine, equine, bovine, rabbit and the like.

**[0038]** The term "alkylparaben" refers to a physiologically tolerated C1-C6 alkyl paraben useful as an antimicrobial agent. Non-limiting examples include at least one methylparaben, ethylparaben, propylparaben, and butylparaben.

**[0039]** The term "aqueous diluent" refers to a liquid solvent that contains water. Aqueous solvent systems may be comprised solely of water, or may be comprised of water plus one or more miscible solvents, and may contain dissolved solutes such as sugars or other excipients. The more commonly-used miscible solvents are the short-chain organic alcohols, such as, methanol, ethanol, propanol, short-chain ketones, such as acetone, and poly Alcohols, such as glycerol.

**[0040]** An "isotonicity agent" is a compound that is physiologically tolerated and imparts a suitable tonicity to a formulation to prevent the net flow of water across cell membranes that are in contact with the formulation. Compounds,

such as glycerin, are commonly used for such purposes at known concentrations. Other suitable isotonicity agents include, but are not limited to, amino acids or proteins (e.g., methionine or albumin), salts (e.g., sodium chloride), and sugars (e.g., dextrose, sucrose and lactose), and/or many others well known in the art, incorporated herein by reference.

**[0041]** The term "preservative" refers to a compound or compositions added to a formulation to act as an anti-microbial, anti-fungal, anti-mycoplasmal, anti-viral, anti-prion and/or anti-bacterial agent. A preserved FSH or FSH variant containing formulation of the present invention preferably meets statutory or regulatory guidelines for preservative effectiveness to be a commercially viable multi-use product. Suitable preservatives can include, but are not limited to, at least one of a benzalkonium chloride, a benzethonium chloride, a chlorohexidine, a phenol, a m-cresol, a benzyl alcohol, an alkyl paraben (methylparaben, ethylparaben, propylparaben, butylparaben, and the like), sodium dehydroacetate, an o-cresol, a p-cresol, a chlorocresol, a phenylmercuric nitrate, a thimerosal, a benzoic acid, and any mixture thereof of one or more preservatives. See, e.g., Wallhauser, K., *Develop. Biol. Standard.* 24, pp. 9-28 (Basel, S. Krager, 1974).

**[0042]** The term "phosphate buffer" refers to excipients that contain a phosphate ion. Generally phosphate buffers are prepared from the phosphoric acid, or salt of phosphoric acid, including but not limited to sodium and potassium salts. Several salts of phosphoric acid are known in the art, such as sodium and potassium monobasic, dibasic, and tribasic salts of the acid. Salts of phosphoric acid are also known to occur as hydrates of the occurring salt. Phosphate buffers may cover a wide range of pHs, such as from about pH 4 to about pH 10, and preferred ranges from about pH 5 to about pH 9, and a most preferred range of about 6.0 to about 8.0. Preferably the formulations of the present invention have pH between about 6.8 and about 7.8, including about pH 7.0, pH 7.2, and 7.4. Preferred ions are the sodium or potassium ions, occurring singularly or together in the solution, as for instance as occurs phosphate buffered saline (PBS). Phosphate saline buffers are well known in the art, such as Dulbecco's Phosphate buffered saline. Salt concentrations in total solution can vary between about 5mM, 9.5mM, 10mM, 50mM, 100mM, 150mM, 200mM, 250mM, and 500mM. Preferably the ion concentration is above 10mM, or above 50mM, or above 100mM.

**[0043]** The term "vial" refers broadly to a reservoir suitable for retaining the FSH and diluent in a contained sterile state. Examples of a vial as used herein include ampules, cartridges, blister packages, or other such reservoir suitable for delivery of the FSH to the patient via pump (including osmotic), catheter, transdermal patch, cartridge, pulmonary, transmucosal, or parenteral delivery. Vials suitable for packaging products for parenteral, pulmonary, transmucosal, or transdermal administration are well-known and recognized in the art.

**[0044]** The term "stability" refers to the physical, chemical, and conformational stability of formulations of FSH of the present invention. Instability of a protein formulation may be caused by chemical degradation or aggregation of the protein molecules to form higher order polymers, by dissociation of the heterodimers into monomers, deglycosylation, modification of glycosylation or any other structural modification that reduces at least one biological activity of an FSH polypeptide included in the present invention.

**[0045]** A "stable" solution or formulation, which is preferable a phosphate buffer with saline or a chosen salt, is one wherein the degree of degradation, modification, aggregation, loss of biological activity and the like, of proteins therein is acceptably controlled, and does not increase unacceptably with time. Stability may be assessed by methods well-known in the art, including measurement of a sample's light scattering, apparent attenuation of light (absorbance, or optical density), size (e.g. by size exclusion chromatography), *in vitro* or *in vivo* biological activity and/or properties by differential scanning calorimetry (DSC). Other methods for assessing stability are well known in the art and can also be used according to the present invention.

**[0046]** The term "treating" refers to the administration, follow up, management and/or care of a patient for which FSH administration is desirable for the purpose of follicle or testicular stimulation or any other physiological response regulated by FSH. Treating can thus include, but is not limited to, the administration of FSH for the induction or improvement of sperm or follicular development or for ovulation induction. In addition, treatments for restoring normal spermatogenesis are contemplated in males.

**[0047]** A "salt" is a physiologically-acceptable salt of FSH. Such salts formed between any one or more of the charged groups in the protein and any one or more physiologically-acceptable, non-toxic cations or anions. Organic and inorganic salts include, for example, those prepared from acids such as hydrochloric, sulfuric, sulfonic, tartaric, fumaric, hydrobromic, glycolic, citric, maleic, phosphoric, succinic, acetic, nitric, benzoic, ascorbic, p-toluenesulfonic, benzenesulfonic, naphthalenesulfonic, propionic, carbonic, and the like, or for example, ammonium, sodium, potassium, calcium, or magnesium. Additional and suitable salts are known in the art and are included herein.

**[0048]** The term "buffer" or "physiologically-acceptable buffer" refers to a compound that is known to be safe for pharmaceutical or veterinary use in formulations and that has the effect of maintaining or controlling the pH of the formulation in the pH range desired for the formulation. Acceptable buffers for controlling pH at a moderately acidic pH to a moderately basic pH include, but are not limited to, such compounds as phosphate, acetate, citrate, arginine, TRIS, and histidine. "TRIS" refers to 2-amino-2-hydroxymethyl-1,3-propanediol, and to any pharmacologically acceptable salt thereof. Preferable buffers are phosphate buffers with saline or an acceptable salt. Other buffers that are physiologically acceptable, and that are suitable for controlling pH at the desired level are known to those of ordinary



skill in the art and are included herein.

**[0049]** A cDNA or genomic library can be screened using a probe based upon the sequence of a polynucleotide or known nucleic acid to obtain a clone encoding a known FSH sequence. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous DNA sequences in the same or different organisms. Those of skill in the art will appreciate that various degrees of stringency of hybridization can be employed in the assay; and either the hybridization or the wash medium can be stringent. As the conditions for hybridization become more stringent, there must be a greater degree of complementarity between the probe and the target for duplex formation to occur. The degree of stringency can be controlled by temperature, ionic strength, pH and the presence of a partially denaturing solvent such as formamide. For example, the stringency of hybridization is conveniently varied by changing the polarity of the reactant solution through, for example, manipulation of the concentration of formamide within the range of 0% to 50%. The degree of complementarity (sequence identity) required for detectable binding will vary in accordance with the stringency of the hybridization medium and/or wash medium. The degree of complementarity will optimally be 100%; however, it should be understood that minor sequence variations in the probes and primers may be compensated for by reducing the stringency of the hybridization and/or wash medium.

**[0050]** Methods of amplification of RNA or DNA are well known in the art and can be used according to the present invention without undue experimentation, based on the teaching and guidance presented herein. Methods of selective amplification by PCR allow for the engineering of smaller segments of nucleic acid sequences, such as those that would encode a defined FSH variant beta chain. Such amplification techniques allow adding convenient termination signals, restrictions sites and the like to the amplified sequence.

**[0051]** Known methods of DNA or RNA amplification include, but are not limited to, polymerase chain reaction (PCR) and related amplification processes (see, e.g., U.S. Patent Nos. 4,683,195, 4,683,202, 4,800,159, 4,965,188, to Mullis, et al.; 4,795,699 and 4,921,794 to Tabor, et al; 5,142,033 to Innis; 5,122,464 to Wilson, et al.; 5,091,310 to Innis; 5,066,584 to Gyllenstein, et al; 4,889,818 to Gelfand, et al; 4,994,370 to Silver, et al; 4,766,067 to Biswas; 4,656,134 to Ringold) and RNA mediated amplification which uses antisense RNA to the target sequence as a template for double-stranded DNA synthesis (U.S. Patent No. 5,130,238 to Malek, et al, with the trade name NASBA), Ausubel, supra; Colligan, supra, Sambrook, supra, the entire contents of which are herein incorporated by reference.

**[0052]** For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of polynucleotides and related DNA sequences directly from genomic DNA or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed as, for example, to obtain any one of the provided FSH or FSH variants, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. Examples of techniques sufficient to direct persons of skill through *in vitro* amplification methods are found in Berger, Sambrook, and Ausubel, supra, as well as Mullis, et al., U.S. Patent No. 4,683,202 (1987); and Innis, et al., PCR Protocols A Guide to Methods and Applications, Eds., Academic Press Inc., San Diego, CA (1990). Commercially available kits for genomic PCR amplification are known in the art. See, e.g., Advantage-GC Genomic PCR Kit (Clontech). The T4 gene 32 protein (Boehringer Mannheim) can be used to improve yield of long PCR products.

**[0053]** Nucleic acids required to express any one of the given FSH or FSH variants can also be prepared by direct chemical synthesis by methods such as the phosphotriester method of Narang, et al., Meth. Enzymol. 68:90-99 (1979); the phosphodiester method of Brown, et al., Meth. Enzymol. 68:109-151 (1979); the diethylphosphoramidite method of Beaucage, et al., Tetra. Letts. 22:1859-1862 (1981); the solid phase phosphoramidite triester method described by Beaucage and Caruthers, Tetra. Letts. 22(20):1859-1862 (1981), e.g., using an automated synthesizer, e.g., as described in Needham-VanDevanter, et al., Nucleic Acids Res. 12:6159-6168 (1984); and the solid support method of U. S. Patent No. 4,458,066. Chemical synthesis generally produces a single-stranded oligonucleotide, which may be converted into double-stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill in the art will recognize that while chemical synthesis of DNA is limited to sequences of about 100 bases, longer sequences may be obtained by the ligation of shorter sequences.

**[0054]** As known in the art one can use recombinant expression cassettes to express known encoding nucleic acids for a known FSH or FSH variant. A nucleic acid sequence, for example a cDNA or a genomic sequence encoding a full-length subunit can be used to construct a recombinant expression cassette which can be introduced into a desired host cell. However, it is appreciated in the art that to obtain functional heterodimers one must express both subunits, whether from one plasmid or introduced on separate plasmids. A recombinant expression cassette will typically comprise a polynucleotide operably linked to transcriptional initiation regulatory sequences that will direct the transcription of the polynucleotide in the intended host cell for each subunit. Such methods are well known in the art to express FSH (e.g. CHO cell-derived recombinant human FSH (rhFSH);(Keene J.L., et al., J. Biol. Chem., 264:4769-4752, 1989; Loumaye E., et al., Human Reprod. Update, 1:188-1999, 1995; Olijve W., et al., Mol. Hum. Reprod., 2:361-369, 1996).

**[0055]** Both heterologous and non-heterologous (i.e., endogenous) promoters can be employed to direct expression of the nucleic acids encoding FSH or FSH variant subunits. General production methodologies by recombinant tech-

niques are well known in the art. See, e.g., Sambrook, et al., 1989; Ausubel, et al., 1987-1989, each entirely incorporated herein by reference.

**[0056]** Encoded polynucleotides for the alpha and beta subunits for FSH or an FSH variant can be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector (or vectors if alpha and beta subunits are contained on separate expression vectors), is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector(s) is a virus, it can be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

**[0057]** The DNA insert for each subunit should be operatively linked to an appropriate promoter, such as the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (e.g., UAA, UGA or UAG) appropriately positioned at the end of the mRNA to be translated.

**[0058]** Expression vector(s) will preferably include at least one selectable marker. Such markers include, e.g., dihydrofolate reductase or neomycin resistance for eucaryotic cell culture, and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal or mammalian cells such as, but not limited to, CHO, COS, AV-12, HEPG2, NIH3T3 and Bowes melanoma cells; and plant cells, with CHO cells preferred. Appropriate culture mediums and conditions for the above-described host cells are known in the art. Preferred eucaryotic vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

**[0059]** Introduction of a vector construct, or vectors, into a host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Sambrook, supra, Chapters 1-4 and 16-18; Ausubel, supra, Chapters 1, 9, 13, 15, 16.

**[0060]** It is anticipated that FSH or a FSH variant subunits can be expressed in a modified form, such as a fusion protein, and can include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, can be added to the N-terminus of a polypeptide to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties can be added to a polypeptide to facilitate purification. Such regions it is anticipated can be removed prior to final preparation of the desired FSH or a FSH variant. Such methods are generally described in many standard laboratory manuals, such as Sambrook, supra, Chapters 17.29-17.42 and 18.1-18.74; Ausubel, supra, Chapters 16, 17 and 18.

**[0061]** Using nucleic acids sequences provided herein or known in the art, one may express the alpha and beta subunits of FSH or a FSH in a recombinantly engineered eucaryotic cell, such as mammalian cells. It is expected that those of skill in the art are knowledgeable in the numerous expression systems available for expression of a nucleic acid encoding a protein that contains two subunits. No attempt to describe in detail the various methods known for the expression of proteins in eucaryotes will be made.

**[0062]** In brief summary, the expression of isolated nucleic acids encoding a known FSH or a FSH variant will typically be achieved by operably linking separately the alpha subunit and the beta subunit DNA or cDNA to a promoter (which is either constitutive or inducible), followed by incorporation into an expression vector(s). Alternatively, by inserting the DNA the vector will provide a suitable promoter. The vector(s) can be suitable for replication and integration in eucaryotic cells. Typical expression vectors contain transcription and translation terminators, initiation sequences and promoters useful for regulation of the expression of the DNA encoding a protein of the present invention. To obtain high level expression of a cloned gene, it is desirable to construct expression vectors which contain, at the minimum, a strong promoter to direct transcription, a ribosome binding site for translational initiation, and a transcription/translation terminator. One of skill in the art would recognize that modifications can be made without diminishing its biological activity. Some modifications may be made to facilitate the cloning, expression, or incorporation of the targeting molecules into the genome. Such modifications are well known to those of skill in the art and include, for example, providing conveniently located restriction sites or termination codons or purification sequences.

**[0063]** Alternatively, nucleic acids can be expressed in a host cell by turning on (by manipulation) in a host cell that contains endogenous DNA encoding the desired alpha and beta subunits. Such methods are well known in the art, e.g., as described in US patent Nos. 5,580,734, 5,641,670, 5,733,746, and 5,733,761, entirely incorporated herein by reference.

**[0064]** A variety of eucaryotic expression systems such as mammalian cells, are known to those of skill in the art. As explained briefly below, a nucleic acid encoding for the alpha and beta subunit of a known FSH or a FSH variant can be expressed in these eucaryotic systems.

**[0065]** Synthesis of heterologous proteins in yeast is well known. F. Sherman, et al., *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory (1982) is a well-recognized work describing the various methods available to produce the protein in yeast. Two widely utilized yeast for production of eucaryotic proteins are *Saccharomyces cerevisiae* and *Pichia pastoris*. Vectors, strains, and protocols for expression in *Saccharomyces* and *Pichia* are known in the art and available from commercial suppliers (e.g., Invitrogen). Suitable vectors usually have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or alcohol oxidase, and an origin of replication, termination sequences and the like as desired.

**[0066]** The sequences encoding the alpha and beta subunits of FSH or a FSH variant can also be ligated to various expression vectors for use in transfecting cell cultures of, for instance, mammalian, insect, or plant origin. Illustrative of cell cultures useful for the production of the peptides are mammalian cells. Mammalian cell systems often will be in the form of monolayers of cells although mammalian cell suspensions may also be used. A number of suitable host cell lines capable of expressing intact proteins have been developed in the art, and include the HEK293, BHK21, and CHO cell lines, with CHO cell lines preferred, such as CHO K1 from Lonza. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter (e.g., preferably the CMV promoter, a HSV tk promoter, EF1 alpha promoter, late or early SV40 promoter, or pgk (phosphoglycerate kinase) promoter), an enhancer (Queen, et al., *Immunol. Rev.* 89:49 (1986)), and processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences. Other animal cells useful for production of proteins of the present invention are available, for instance, from the American Type Culture Collection Catalogue of Cell Lines and Hybridomas (7th edition, 1992). Preferred host cells include CHO cells, such as CHO-K1 and preferred vectors include GS vectors, each available, e.g., from Lonza Biologics PLC (Slough, Berkshire, England, UK).

**[0067]** Appropriate vectors for expressing the alpha and beta subunit of FSH or a FSH variant in insect cells are usually derived from the SF9 baculovirus. Suitable insect cell lines include mosquito larvae, silkworm, armyworm, moth and *Drosophila* cell lines such as a Schneider cell line (See Schneider, J. *Embryol. Exp. Morphol.* 27:353-365 (1987)).

**[0068]** As with yeast, when higher animal or plant host cells are employed, polyadenylation or transcription terminator sequences are typically incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript may also be included. An example of a splicing sequence is the VP1 intron from SV40 (Sprague, et al., *J. Virol.* 45:773-781 (1983)). Additionally, gene sequences to control replication in the host cell may be incorporated into the vector such as those found in bovine papilloma virus type-vectors. M. Saveria-Campo, *Bovine Papilloma Virus DNA*, a Eucaryotic Cloning Vector in *DNA Cloning Vol. II*, a Practical Approach, D. M. Glover, Ed., IRL Press, Arlington, VA, pp. 213-238 (1985).

**[0069]** FSH or a FSH variant, once expressed, can be isolated from the cells by applying standard protein isolation techniques to the lysates. The monitoring of the purification process can be accomplished by using Western blot techniques or radioimmunoassay of other standard immunoassay techniques.

**[0070]** FSH or a FSH variant, containing an alpha and beta subunit, can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, size exclusion chromatography, and lectin chromatography. Preferably, high performance liquid chromatography ("HPLC"), cation exchange chromatography, affinity chromatography, size exclusion chromatography, or combinations thereof, are employed for purification. FSH and FSH variants having an alpha and beta subunit include naturally purified products, products of chemical synthetic procedures, and include products produced by recombinant techniques from a eucaryotic host, including, for example, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention can be glycosylated or can be non-glycosylated. Preferred FSH or a FSH variant molecules are glycosylated as would occur in eucaryotic hosts. In addition, polypeptides of the invention can also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Such methods are described in many standard laboratory manuals, such as Sambrook, supra, Chapters 17.37-17.42; Ausubel, supra, Chapters 10, 12, 13, 16, 18 and 20, entirely incorporated herein by reference.

**[0071]** FSH or a FSH variant known in the art, include but are not limited to those protein sequences listed in the sequence identification portion of the specification of which are further identified below:

SEQ ID NO: 1;	bovine alpha subunit	- 96 amino acids
SEQ ID NO: 2;	bovine beta subunit	- 111 amino acids
SEQ ID NO: 3;	equine alpha subunit	- 96 amino acids
SEQ ID NO: 4;	equine beta subunit	- 111 amino acids
SEQ ID NO: 5;	human alpha subunit	- 92 amino acids
SEQ ID NO: 6;	human beta subunit	- 111 amino acids

(continued)

SEQ ID NO: 7;	porcine alpha subunit	- 96 amino acids
SEQ ID NO: 8;	porcine beta subunit	- 111 amino acids
SEQ ID NO: 9;	ovine alpha subunit	- 96 amino acids
SEQ ID NO: 10;	ovine beta subunit	- 111 amino acids-
SEQ ID NO: 11;	human beta variant	- 108 amino acids
SEQ ID NO: 12;	human beta variant	- 109 amino acids
SEQ ID NO: 13;	human beta variant	- 110 amino acids

**[0072]** FSH or a FSH variant nucleotide sequence, include but are not limited to those nucleotide sequences which encode an alpha or a beta subunit listed in the sequence identification portion of the specification of which are further identified below:

SEQ ID NO:14;	human alpha cDNA (codes 92 amino acids)	- 276 nucleotides
SEQ ID NO:15;	h. beta variant cDNA (codes 108 amino acids)	- 324 nucleotides
SEQ ID NO:16;	h. beta variant cDNA (codes 109 amino acids)	- 327 nucleotides
SEQ ID NO:17;	h. beta variant cDNA (codes 110 amino acids)	- 330 nucleotides
SEQ ID NO:18;	h. beta cDNA (codes 111 amino acids)	- 333 nucleotides
SEQ ID NO:19;	human alpha cDNA (codes 92 amino acids)	- 276 nucleotides
SEQ ID NO:20;	h. beta variant cDNA (codes 108 amino acids)	- 324 nucleotides

The DNA of SEQ ID NO:19 and 20 is designed and constructed from ligated oligonucleotides. The differences between SEQ ID NO:19 and SEQ ID NO:14 are one that do not change the encoded amino acid sequence of the alpha subunit protein. Likewise, the differences between SEQ ID NO:20 and SEQ ID NO:15 are ones that do not change the encoded amino acid sequence of the beta variant subunit protein.

**[0073]** The amino acids that make up the proteins and polypeptides of the present invention are often abbreviated. The amino acid designations can be indicated by designating the amino acid by its single letter code, its three letter code, name, or three nucleotide codon(s) as is well understood in the art (see Alberts, B., et al., Molecular Biology of The Cell, Third Ed., Garland Publishing, Inc., New York, 1994):

SINGLE LETTER CODE	THREE LETTER CODE	NAME	THREE NUCLEOTIDE CODON(S)
A	Ala	Alanine	GCA, GCC, GCG, GCU
C	Cys	Cysteine	UGC, UGU
D	Asp	Aspartic acid	GAC, GAU
E	Glu	Glutamic acid	GAA, GAS
F	Phe	Phenylalanine	UUC, UUU
G	Gly	Glycine	GGA, GGC, GGG, GGU
H	His	Histidine	CAC, CAU
I	Ile	isoleucine	AUA, AUC, AUU
K	Lys	Lysine	AAA, AAG
L	Leu	Leucine	UUA, UUG, CUA, CUC, CUG, CUU
M	Met	Methionine	AUG

(continued)

SINGLE LETTER CODE	THREE LETTER CODE	NAME	THREE NUCLEOTIDE CODON(S)
N	Asn	Asparagine	AAC, AAU
P	Pro	Proline	CCA, CCC, CCG, CCU
Q	Gln	Glutamine	CAA, CAG
R	Arg	Arginine	AGA, AGG, CGA, CGC, CGG, CGU
S	Ser	Serine	AGC, AGU, UCA, UCC, UCG, UCU
T	Thr	Threonine	ACA, ACC, ACG, ACU
V	Val	Valine	GUA, GUC, GUG, GUU
W	Trp	Tryptophan	UGG
Y	Tyr	Tyrosine	UAC, UAU

[0074] As noted above, the invention provides for stable formulations, which is preferable a phosphate buffer with saline or a chosen salt, as well as preserved solutions and formulations containing a preservative as well as multi-use preserved formulations suitable for pharmaceutical or veterinary use, comprising FSH or FSH variant in a pharmaceutically acceptable formulation. Preserved formulations contain at least one preservative selected from the group consisting of at least one phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent.

[0075] As noted above, the invention provides an article of manufacture, comprising packaging material and a vial comprising a solution of FSH or a FSH variant with the prescribed buffers and/or preservatives, optionally in an aqueous diluent, wherein said packaging material comprises a label which indicates that such solution may be held over a period of twenty-four hours or greater. The invention further comprises an article of manufacture, comprising packaging material, a first vial comprising lyophilized FSH or a FSH variant, and a second vial comprising an aqueous diluent of prescribed buffer or preservative, wherein said packaging material comprises a label which instructs a patient to reconstitute the FSH or a FSH variant in the aqueous diluent to form a solution which may be held over a period of twenty-four hours or greater.

[0076] The FSH or a FSH variant used in accordance with the present invention may be produced by recombinant means, including from mammalian cell or transgenic preparations, or may be purified from other biological sources, such as from urinary sources. Acceptable methodologies include those described in Hakola, K. Molecular and Cellular Endocrinology, 127:59-69, 1997; Keene, et al., J. Biol. Chem., 264:4769-4775, 1989; Cerpa-Poljak, et al., Endocrinology, 132:351-356, 1993; Dias, et al., J. Biol. Chem., 269:25289-25294, 1994; Flack, et al., J. Biol. Chem., 269:14015-14020, 1994; and Valove, et al., Endocrinology, 135:2657-2661, 1994, and U.S. Patent 3,119,740, herein entirely incorporated by reference.

[0077] The method by which the proteins are provided for the formulations of this invention is not particularly relevant. Preferably FSH is a heterodimer comprising one alpha subunit and one beta subunit, respectfully, as provide in SEQ ID NOS 5 and 6, or a FSH variant heterodimer comprising one alpha subunit and one beta subunit, respectfully, as given in SEQ ID NOS: 5 and 11; 5 and 12; and 5 and 13. Suitable FSH or a FSH variant species within the present invention include, but are not limited to, at least one known alpha subunit sequence and at least one known beta subunit (see sequence listing for known alpha and beta subunits and as otherwise known in the art).

[0078] Non-limiting examples of FSH or a FSH variant, include but are not limited to:

(a) : $\alpha$ -subunit: (SEQ ID NO:1)

FPDGEFTMQGCPECKLKENKYFSKPDAPYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:2)

RSCELTNITITVEKEECGFCISINTWCAGYCYTRDLVYRDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCSKCDSDSTDCTVRGLGPSYCSFREIKE

(b): $\alpha$ -subunit:(SEQ ID NO:3)

FPDGEFTTQDCPECKLRENKYFFKLGVPPIYQCKGCCFSRAYPTPARSRKTM LVPKN  
ITSESTCCVAKAFIRVTVMGNIKENHTQCYCSTCYHHKI

$\beta$ -subunit:(SEQ ID NO:4)

NSCELTNITIAVEKEGCGFCITINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATACHCGKCNSDSTDCTVRGLGPSYCSFGDMKE

(d): $\alpha$ -subunit:(SEQ ID NO:7)

(c): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:6)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMKE

FPDGEFTMQGCPECKLKENKYFSKLGAPIYQCMGCCFSRAYPTPARSKKTMLV PKN  
ITSEATCCVAKAFTKATVMGNARVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:8)

NSCELTNITITVEKEECNFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCGKCDSDSTDCTVRGLGPSYCSFSEMKE

(e): $\alpha$ -subunit:(SEQ ID NO:9)

FPDGEFTMQGCPECKLKENKYFSKPDAPIYQCMGCCFSRAYPTPARSKKTMLV PKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:10)

RSCELTNITITVEKEEC SFCISINTTWCAGYCYTRDLVYKDPARPNIQKACTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCGKCDRDSTDCTVRGLGPSYCSFSDIRE

(f): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:11)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGE

(g) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:12)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEM

(h) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:13)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMK

The range of protein hormone in the product of the present invention includes amounts yielding upon reconstitution, if in a wet/dry system, concentrations from about 1.0  $\mu$ g/ml to about 50 mg/ml, although lower and higher concentrations are operable and are dependent on the intended delivery vehicle, e.g., solution formulations will differ from transdermal patch, pulmonary, transmucosal, or osmotic or micro pump methods. The hormone concentrations are preferably about 5.0  $\mu$ g/ml to 2 mg/ml and most preferably about 5.0  $\mu$ g/ml, or 10  $\mu$ g/ml, or 50  $\mu$ g/ml to about 200  $\mu$ g/ml.

**[0079]** Preferably, the aqueous diluent optionally further comprises a pharmaceutically acceptable preservative. Preferred preservatives include those selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof. Preferably, the preservative is meta-cresol, phenol, chlorocresol, or a mixture thereof, with m-cresol most preferred. The concentration of preservative used in the formulation is a concentration sufficient to yield an anti-microbial effect. Such concentrations are dependent on the preservative selected and are readily determined by the skilled artisan. For example, m-cresol or phenol (alone or in combination) are generally at a concentration from about 23 mM to about 35 mM. Surprisingly, the preservatives used in the presently claimed formulations do not adversely affect the biological activity of FSH or a FSH variant and allow multi-use administration.

**[0080]** Other excipients, e.g. isotonicity agents, buffers, antioxidants, preservative enhancers, may be optionally and preferably added to the diluent. An isotonicity agent, such as glycerin, is commonly used at known concentrations. The concentration of glycerin is generally about 16 mg/ml. A physiologically tolerated buffer is preferably added to provide improved pH control. The formulations may cover a wide range of pHs, such as from about pH 4 to about pH 10, and preferred ranges from about pH 5 to about pH 9, and a most preferred range of about 6.0 to about 8.0. Preferably the formulations of the present invention have pH between about 6.8 and about 7.8. Preferred buffers include phosphate buffers, most preferably sodium phosphate, particularly phosphate buffered saline (PBS).

**[0081]** Other additives, such as a pharmaceutically acceptable solubilizers like Tween 20 (polyoxyethylene (20) sorbitan monolaurate), Tween 40 (polyoxyethylene (20) sorbitan monopalmitate), Tween 80 (polyoxyethylene (20) sorbitan monooleate), Pluronic F68 (polyoxyethylene polyoxypropylene block copolymers), and PEG (polyethylene glycol) or non-ionic surfactants such as polysorbate 20 or 80 or poloxamer 184 or 188, Pluronic® polyols, other block co-polymers, and chelators such as EDTA and EGTA may optionally be added to the formulations or compositions to reduce aggregation. These additives are particularly useful if a pump or plastic container is used to administer the formulation. The presence of pharmaceutically acceptable surfactant mitigates the propensity for the protein to aggregate. The present claimed formulations are surprisingly stable. Prior to the present invention, the preparation of preserved, multi-use formulations of FSH was believed to be impossible due to instability. Applicants have discovered that the claimed formulations may be safely stored at temperatures of from about 2 to about 40°C and retain the biological activity of the protein for extended periods of time, exceeding 2 months and as further demonstrated.

**[0082]** The formulations of the present invention can be prepared by a process which comprises mixing FSH or a FSH variant and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben, (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride,

sodium dehydroacetate and thimerosal or mixtures thereof in an aqueous diluent. Mixing the FSH or a FSH variant and preservative in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of FSH or a FSH variant in buffered solution is combined with the desired preservative in a buffered solution in quantities sufficient to provide the protein and preservative at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimized for the concentration and means of administration used.

**[0083]** The claimed formulations may be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized FSH or a FSH variant that is reconstituted with a second vial containing a preservative and/or excipients, preferably a phosphate buffer and/or saline and a chosen salt, in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution may be reused multiple times and may suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

**[0084]** The present claimed articles of manufacture are surprisingly useful for administration over a period of twenty-four hours or greater. Prior to the present invention, such products were only suitable and approved for immediate use. The patient was instructed to discard unused material leading to waste and expense. Accordingly, the presently claimed articles of manufacture offer significant advantages to the patient. Applicants have discovered that the claimed formulations may be safely stored at temperatures of from about 2 to about 40°C and retain the biological activity of the protein for extended periods of time, exceeding 2 months; thus, allowing a package label indicating that the solution may be held and/or used over a period of 24, 36, 48, 72, or 96 hours or greater. If preserved diluent is used, such label may include use up to one, one and a half, to two years.

**[0085]** The solutions of FSH or a FSH variant in the invention can be prepared by a process which comprises mixing FSH or a FSH variant in an aqueous diluent. Mixing is carried out using conventional dissolution and mixing procedures. To prepare a suitable diluent, for example, a measured amount of FSH or a FSH variant in water or buffer is combined in quantities sufficient to provide the protein and optionally a preservative or buffer at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimized for the concentration and means of administration used.

**[0086]** The claimed products may be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized FSH or a FSH variant that is reconstituted with a second vial containing the aqueous diluent. Either a single solution vial or dual vial requiring reconstitution may be reused multiple times and may suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

**[0087]** The claimed products may be provided indirectly to patients by providing to pharmacies, clinics, or other such institutions and facilities, clear solutions or dual vials comprising a vial of lyophilized FSH or a FSH variant that is reconstituted with a second vial containing the aqueous diluent. The clear solution in this case may be up to one liter or even larger in size, providing a large reservoir from which smaller portions of the FSH or a FSH variant solution may be retrieved one or multiple times for transfer into smaller vials and provided by the pharmacy or clinic to their customers and/or patients. The diluent vial in this case may be up to one liter or even larger in size, providing a large reservoir from which smaller portions of the diluent may be retrieved multiple times for reconstitution of the lyophilized FSH or a FSH variant. The clear solution or reconstituted FSH or a FSH variant solution provided by the pharmacy or clinic to their customers and patients may suffice for single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

**[0088]** Recognized devices comprising these single vial systems include those pen-injector devices for delivery of a solution such as Humaject®, NovoPen®, B-D®Pen, AutoPen®, and OptiPen®. Recognized devices comprising a dual vial system include those pen-injector systems for reconstituting a lyophilized drug in a cartridge for delivery of the reconstituted solution such as the HumatroPen.

**[0089]** The products presently claimed include packaging material. The packaging material provides, in addition to the information required by the regulatory agencies, the conditions under which the product may be used. The packaging material of the present invention provides instructions to the patient to reconstitute the FSH or a FSH variant in the aqueous diluent to form a solution and to use the solution over a period of twenty-four hours or greater for the two vial, wet/dry, product. For the single vial, solution product, the label indicates that such solution may be used over a period of twenty-four hours or greater. The presently claimed products are useful for human pharmaceutical product use.

**[0090]** The formulations of the present invention can be prepared by a process which comprises mixing FSH or a FSH variant and a selected buffer, preferably a phosphate buffer containing saline or a chosen salt. Mixing the FSH or a FSH variant and buffer in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of FSH or a FSH variant in water or buffer is combined with the desired buffering agent in water in quantities sufficient to provide the protein and buffer at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order



the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimized for the concentration and means of administration used.

**[0091]** The claimed stable or preserved formulations may be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized FSH or a FSH variant that is reconstituted with a second vial containing a preservative or buffer and excipients in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution may be reused multiple times and may suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

**[0092]** FSH or a FSH variant in either the stable or preserved formulations or solutions described herein, may be administered to a patient in accordance with the present invention via a variety of delivery methods including SC or IM injection; transdermal, pulmonary, transmucosal, implant, osmotic pump, cartridge, micro pump, oral, or other means appreciated by the skilled artisan, as well-known in the art.

**[0093]** The following examples are provided merely to further illustrate the preparation of the formulations and compositions of the invention. The scope of the invention shall not be construed as merely consisting of the following examples.

#### Example 1

##### Cloning and Expression of FSH or a FSH variant, in Mammalian Cells

**[0094]** A typical mammalian expression vector contains at least one promoter element, which mediates the initiation of transcription of mRNA, the polypeptide coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. However, because functional FSH or FSH variants contain both an alpha and beta subunit, means to express both subunits are required, either by expressing both subunits from single vector containing a promoter element for each subunit, or by using two vectors: a first vector containing a promoter to express the first subunit and a second vector that has a promoter to express the second subunit.

**[0095]** Additionally, each mammalian expression vector have elements that may be present on one or more vectors include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing.

**[0096]** Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in providing FSH or FSH variants subunits include, for example, vectors such as pRES1neo, pRetro-Off, pRetro-On, PLXSN, or pLNCX (Clontech Labs, Palo Alto, CA), pcDNA3.1 (+/-), pcDNA/Zeo (+/-) or pcDNA3.1/Hygro (+/-) (Invitrogen), PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include human Hela 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

**[0097]** Alternatively, the desired DNA sequences for the alpha and beta subunit can be expressed in stable cell lines that contain the DNA sequences for expressing each subunit once integrated into a chromosome(s). The co-transfection with a selectable marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells as known in the art.

**[0098]** The transfected DNA sequences for the subunits can also be amplified to express large amounts of the encoded polypeptide. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the DNA sequences of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy, et al., Biochem. J. 227:277-279 (1991); Bebbington, et al., Bio/Technology 10: 169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins and polypeptides.

**[0099]** The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen, et al., Molec. Cell. Biol. 5:438-447 (1985)) plus a fragment of the CMV-enhancer (Boshart, et al., Cell 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the DNA sequences for the alpha and beta subunits of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

#### Example 2

##### Cloning and Expression in COS or CHO Cells

**[0100]** An expression plasmid for FSH or a FSH variant is made by cloning a cDNA encoding FSH or a FSH variant

subunits into the expression vector pcDNAI/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.). As previously mentioned, each subunit requires expression to produce a function heterodimer, either from independent introduction of separate vectors into the host cell or by engineering a single vector to express both alpha and beta subunits.

**[0101]** The expression vector(s) pcDNAI/amp contains: (1) an E. coli origin of replication effective for propagation in E. coli and other prokaryotic cells; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eucaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron; (5) several codons encoding a hemagglutinin fragment (i.e., an "HA" tag to facilitate purification) or HIS tag (see, e.g., Ausubel, supra) followed by a termination codon and polyadenylation signal arranged so that a cDNA can be conveniently placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker. The HA tag corresponds to an epitope derived from the influenza hemagglutinin polypeptide described by Wilson, et al., Cell 37:767-778 (1984). The fusion of the HA tag to the target polypeptide, either the alpha or beta subunit, allows easy detection and recovery of the recombinant polypeptide with an antibody that recognizes the HA epitope. pcDNAIII contains, in addition, the selectable neomycin marker.

**[0102]** A DNA fragment encoding the alpha and beta subunit of FSH or a FSH variant, is separately cloned into the polylinker region of the vector so that recombinant polypeptide expression is directed by the CMV promoter. Insertion into the vector is optionally with or without the HA epitope. The plasmid construction strategy is as follows. The FSH or a FSH variant, cDNA of the deposited clone for each subunit is amplified using primers that contain convenient restriction sites.

**[0103]** The PCR amplified DNA fragment for each subunit and the vector, pcDNAI/Amp, are digested with suitable restriction enzyme(s) and then each subunit is ligated to digested vector. Each ligation mixture is transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037), and the transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA for each subunit is isolated from resistant colonies and examined by restriction analysis or other means for the presence of the FSH or a FSH variant encoding fragment.

**[0104]** For expression of recombinant FSH or a FSH variant, COS cells are co-transfected with an expression vector for each subunit, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook, et al., Molecular Cloning: a Laboratory Manual, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989). Cells are incubated under conditions for expression of FSH or a FSH variant, by each vector.

**[0105]** It is expected that expression of the FSH HA or FSH variant HA fusion polypeptide is detected by radiolabeling and immunoprecipitation, using methods described in, for example Harlow, et al., Antibodies: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). To this end, two days after transfection, the cells are labeled by incubation in media containing 35S-cysteine for 8 hours. The cells and the media are collected, and the cells are washed and lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson, et al. cited above. Proteins are precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated protein is then analyzed by SDS-PAGE and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

**[0106]** The vector pC4 is used for the expression of each subunit of FSH or a FSH variant. Alternatively, one skilled in the art would be able to adapt pC4 to express both alpha and beta subunits from a single vector. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary or other cells lacking dihydrofolate activity that are co-transfected with alpha and beta subunit plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., F. W. Alt, et al., J. Biol. Chem. 253:1357-1370 (1978); J. L. Hamlin and C. Ma, Biochem. et Biophys. Acta 1097:107-143 (1990); and M. J. Page and M. A. Sydenham, Biotechnology 9:64-68 (1991)). Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If DNA sequences are linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach can be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained which contain the amplified DNA sequences integrated into one or more chromosome(s) of the host cell.

**[0107]** Plasmid pC4 contains for expressing the alpha and beta subunit DNA sequences of interest behind the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen, et al., Molec. Cell. Biol. 5:438-447 (1985)) which additionally contains a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart, et al., Cell 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the DNA sequences. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters

can also be used for the expression, e.g., the human b-actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the FSH or a FSH variant, in a regulated way in mammalian cells (M. Gossen, and H. Bujard, Proc. Natl. Acad. Sci. USA 89: 5547-5551 (1992)). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying the DNA sequences of the alpha and beta subunit integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

**[0108]** The plasmid pC4 is digested with restriction enzymes and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

**[0109]** The DNA sequence encoding the complete FSH or a FSH variant, for each subunit is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. Non-limiting examples include 5' and 3' primers having nucleotides corresponding or complementary to a portion of the coding sequence of each subunit for an FSH or a FSH variant according to known methods in the art.

**[0110]** The amplified fragment(s) are digested with suitable endonucleases and then purified again on a 1% agarose gel. The isolated fragments for each subunit and the dephosphorylated vector are then separately ligated with T4 DNA ligase. E. coli HB101 or XL-1 Blue cells are then separately transformed and bacteria are identified that contain the fragment (or fragments if the vector is adapted for expressing both alpha and beta subunits) inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

**[0111]** Chinese hamster ovary (CHO) cells lacking an active DHFR gene are used for transfection. 5 µg of the expression plasmid(s) pC4 is cotransfected with 0.5 µg of the plasmid pSV2-neo using lipofectin. The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 µg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 µg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 mM, 2 mM, 5 mM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 mM. Expression of the desired product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

**[0112]** It will be clear that the invention can be practiced otherwise than as particularly described in the foregoing description and examples.

**[0113]** Numerous modifications and variations of the present methodologies is known in the art to expressing recombinant proteins, including those as described in Keene J.L., et al., J. Biol. Chem., 264:4769-4752, 1989; Loumaye E., et al., Human Reprod. Update, 1:188-1999, 1995; Olijve W., et al., Mol. Hum. Reprod., 2:361-369, 1996, as well as other recombinant techniques known and used for other gonadotropins.

### Example 3

#### Expression in AV12 Cells

**[0114]** An expression cassette vector pGTH was used for expression of the alpha subunit in AV12. A number of published articles describe the use of AV12-664 and/or AV12-RGT18 cells (see Grinnell, B.W. et al., Blood 76(12):2546-54, 1990; Burck, P.J. et al., J. Biol. Chem. 265(9):5170-7, 1990; Parkinson, J.F. et al., J. of Biol. Chem. 265(21):12602-10, 1990; Grinnell, B.W. et al., J. Biol. Chem 266(15):9778-85, 1991; Wery, J.P. et al., Nature 352(6330):79-82, 1991; Berg, D.T. et al., Biotechniques 14:972-9, 1993; Gerlitz, B. et al. Biochemical Journal 295 (1):131-40, 1993; Kursar, J.D. et al. Molecular Pharmacology 46(2):227-34, 1994; Desai, M.A. et al. Molecular Pharmacology 48(4):648-57, 1995; Obesity Research 3 Suppl 4:4441S-4447S, 1995; Desai, M.A. et al. British Journal of Pharmacology 118(6):1558-64, 1996; Kumar, A. et al. Cancer Research 56(23):5397-402, 1996; Boggs, L.N. et al. J. of Neurochemistry 57(3):1324-7, 1996; Lucaites, V.L. et al. Life Sciences 59(13):1081-95, 1996; Schoepp, D.D. et al. Neuropharmacology 35(12):1661-72, 1996; Kumar, A. et al. Cancer Research 57(15):3111-4, 1997; Urology 49(3):487-93, 1997; Wainscott, D.B. et al. Naunyn-Schmiedeberg's Archives of Pharmacology 357(1):17-24, 1998; Wu, S. et al. Brain Research - Molecular Brain Research 53(1-2):88-97, 1998. Similarly, publications have described use of the plasmid pGTH (see Grinnell, et al. J. Biol. Chem. 266(15):9778-85, 1991) and the plasmid pGTD (see Biochemical Journal 295 (Pt1): 131-40, 1993).

**[0115]** Briefly, pGTH contains sequentially several elements: the SV40 early promoter, ori, E.coli hygromycin resistance, SV40 small "t" antigen splice site/poly-A site, pBR322 cloning remnant, BK virus (strain P2) cloning remnant, Poly-CA<sub>20</sub>/GT<sub>20</sub> element (synthetic oligonucleotide), BK virus (strain P2) enhancer, AD2 major late promoter/spliced

tripartite leader, BclI insertion site for the FSH alpha subunit coding sequence (including stop codon), SV40 small "t" antigen splice site/poly-A site; and pBR322 ampicillin resistance/ori.

**[0116]** The plasmid construct pGTH-alpha was generated to express the encoded the human alpha subunit sequence (SEQ ID:5) by cloning a 362-bp BclI FSH cDNA fragment into the unique BclI site of the vector (see sequence - SEQ ID:19 or 14). The FSH alpha cDNA fragment DNA was generated by PCR amplification using the shuttle plasmid pLGD637 as template (pLGD637 contains a synthetic/oligonucleotide-assembled FSH alpha cDNA sequence). The integrity of BclI insert was confirmed by sequencing followed by comparison to the GenPept database (Accession Number 31869).

**[0117]** An expression cassette vector pGTD was used to express the human beta subunit FSH variant sequence (SEQ ID:11). pGTD contains several elements for expression in AV12 cells. pGTD contains sequentially the BK virus (strain P2) cloning remnant, Poly-CA<sub>20</sub>/GT<sub>20</sub> element (synthetic oligonucleotide), BK virus (strain P2) enhancer, AD2 major late promoter/spliced tripartite leader, BclI insertion site for the FSH variant beta subunit coding sequence (including stop codon), SV40 small "t" antigen splice site/poly-A site; SV40 early promoter.ori, Murine dihydrofolate reductase cDNA, SV40 small "t" antigen splice site/poly-A site, and pBR322 ampicillin resistance/ori.

**[0118]** The plasmid construct pGTD-bCD3 was generated by cloning a 393-bp BclI FSH beta-bCD3 cDNA fragment into the unique BclI site of the pGTD vector (see SEQ ID:20 or 15). The FSH beta-CD3 cDNA fragment DNA was generated by PCR amplification, using the shuttle plasmid pLGD638 as template (pLGD638 contains a synthetic/oligonucleotide-assembled FSH beta cDNA sequence). The integrity of the construct was confirmed by sequencing and compared with the human beta subunit sequence deposited in the GenPept database (Accession Number 476441).

**[0119]** In brief, the pGTH-alpha and pGTD-bCD3 plasmids were linearized, repurified, and then co-transfected into adherent AV23-RGT18 cells. Following selection with medium containing 0.25uM methotrexate and 100 µg/ml hygromycin-B, along with 200 µg/ml G418 to maintain the glutamate transporter genotype of the AV12-RGT18 cells, individual stable clones were isolated either manually or via flow-assisted cell sorting. Highest producing clones were identified by analysis of conditioned medium with a commercial FSH Elisa kit. Several clones were adapted to serum-free suspension and further amplified to obtain isolatable quantities of the FSH variant heterodimer.

#### Example 4

##### Expression In CHO-K1 Cells

**[0120]** A CHO-K1 cell line (LONZA Biologics plc.) was developed to produce a FSH variant heterodimer composed of the alpha subunit of SEQ ID NO:5 and the beta subunit of SEQ ID NO:11.

**[0121]** The expression vector cassette contains the encoding DNA for alpha subunit, SEQ ID NO:5, and the encoding DNA for the beta subunit, SEQ ID NO:11. are controlled by two different promoters: CMV for the beta subunit and EF1 Alpha for the alpha subunit. Each alpha and beta subunit sequence uses the Bovine Growth Hormone polyA tail. Additionally the vector contains the Glutamine Synthetase gene, controlled by the SV40 Late Promoter and containing the SV40 polyA tail, is used as the selectable marker. This vector was used to transfect the CHO-K1 cells.

**[0122]** The cell line was grown in GibcoBRL's CD CHO media under selective pressure of L-methionine sulfoximine. ELISA assays were used to identify master wells expressing FSH variant. Several master wells were subjected to cloning and amplification procedures. These experiments led to the clonal cell line 2B6.1C3.25 which had suitable titers. Expression studies conducted in small scale shake flasks (20-60 ml) have shown that this line expresses FSH variant at 30 mg/L after 7 to 8 days.

#### Example 5

##### Purification of FSH variants from CHO or AV-12 Cells

**[0123]** Purification of FSH variant heterodimer comprised of an alpha subunit of SEQ ID NO:5 and a beta subunit of SEQ ID:11 can be accomplished by a number of methods described and known in the art from monolayer or suspension cultures of either CHO-K1 or AV12 cell lines or other production lines suitably available. One method for isolating the disclosed FSH variant from the culture containing medium is subjecting the culture medium to cation exchange chromatography, dye affinity chromatography and gel filtration chromatography to purify the protein. In the case of suspension cultures, which may contain detergents, additional purification steps may be needed such as a Q-Sepharose step. The chromatographic steps can be further added to or optimized for pH, conductivity, buffer composition and running conditions (column dimensions, flow rates, etc). Purity and yield can be analyzed by SDS-PAGE gels (both Coomassie staining and Western blotting), ELISA assays, exclusion chromatography and protein concentration by UV absorbance at 277nm or other known techniques.

**[0124]** Purification and chromatographic fractionation can be achieved by following the further details for each iso-

lation step given below. For monolayer cultures, the conditioned medium is concentrated and diafiltered prior to application to the cation exchange column. The Q-Sepharose batching step can be included for suspension cultures to remove detergents that may be present.

## 1. Concentration

**[0125]** Typically if 0.02 to 0.04% Pluronic F68 is used for AV12 suspension cultures, whereas there is 0.1 to 0.18% in the media used for CHO suspension cultures. The conditioned medium, clarified by the cell culture group using centrifugation and filtration through a cheese cloth, is concentrated using a tangential flow filtration system ProFlux (ProFlux M12 from Millipore) with a S3YM10 spiral cartridge (Amicon #540633). Depending on the amount of Pluronic F68 that was used, the medium is concentrated 4 to 10-fold so that there is 0.2-0.4% Pluronic F68 in the concentrated medium. The final volume of material after concentration is usually 2-3L starting from 8L for CHO-K1 and 24L for AV12 cultures.

## 2. Batching with Q-Sepharose

**[0126]** For each liter of starting conditioned medium before concentration, 50 ml of Fast Flow Q-Sepharose resin (Pharmacia 17-0510-01, pre-equilibrated with 20 mM Tris, pH 7.4) and sufficient NaCl to give a final conductivity of 200 mM (~20 mS/cm) are added to the concentrated medium and stirred gently overnight at 4°C

## 3. Diafiltration

**[0127]** After overnight batching of the concentrated medium with Q-Sepharose resin, the resin is allowed to settle, the supernatant is decanted out and filtered using a CUNO system with a Zeta Plus 30-SP Filter (#B0406-30SP from Sun Source Fauver) and a Masterflex pump at a flow rate of 170 ml/min. The medium is then further concentrated to about 800 ml and diafiltered using 5-6 volumes of 20 mM Tris pH 7.4 in the Proflux system. At this point, the conductivity is ~ 2 mS/cm. The pH is adjusted to 5.0 with 1N HCl and the solution is again filtered using a fresh Zeta Plus 30-SP filter in the CUNO system and immediately loaded onto the Cation Exchange column.

## 4. Cation Exchange Chromatography (CEX)

**[0128]** Column: Pharmacia SP-Sepharose Fast Flow (17-0729-01) is used to pack a 50 ml column for ~100-200 mg of FSH (total protein ~ 500-600 mg). Buffers are: A: 20 mM sodium phosphate, pH 5.0; and B: 20 mM sodium phosphate, pH 5.0, 1M NaCl

**[0129]** The sample is adjusted to pH 5.0, clarified by filtration and immediately loaded onto the column and run at 5 ml/min with 4 Column Volumes before starting a gradient of 0% to 50% B over 15 Column Volumes. 3 min fractions are collected (15 ml). Fractions are collected into 400 ml of 1 M Tris, pH 8.0.

**[0130]** Coomassie-stained SDS PAGE gels are used to choose FSH-containing fractions to pool (typically the pool size is 200 to 250 ml for a 50 ml column). This pool is dialyzed against 20 mM Tris, pH 7.4 overnight at 4°C to drop conductivity to < 3 mS/cm.

## 5. Dye Affinity Chromatography (DAC)

**[0131]** Column: Mimetic Blue Dye 1 A6XL, 0090-0500 from Prometic Biosciences Inc. is used in a 50 ml column for ~40 mg of FSH. Buffers are: A: 25 mM phosphate, pH 6.5 (conductivity is 4.5 to 5 mS/cm); B: 25 mM phosphate, 150 mM KCl, pH 6.5; C: 25 mM phosphate, 1M KCl, pH 8.0

**[0132]** After dialysis of CEX pool, the pH is adjusted to 6.5, and loaded onto the DAC column at 3 ml/min. A gradient from 100%A to 50%A; 50% B is applied for 4 Column Volumes, then eluted with 100% Buffer C for 5 Column Volume, collecting Fractions of 3 min (9 ml).

**[0133]** Coomassie-stained SDS-PAGE gels are used to choose FSH-containing fractions to pool. Typically, the pool size is 90 to 100 ml for a 50 ml column. The pool is concentrated to 4 ml using Millipore Ultrafree centrifuge devices (UFV2BCC40, 5000 MWCO, spun at 2000 rpm) and loaded onto a Gel Filtration column.

## 6. Gel Filtration

**[0134]** Column: BioPilot Superdex 75 Prep Grade 35/600 column is used for 50 to 100 mg of FSH. Buffers are: 1x PBS (made from GIBCO 10x PBS, #70011) plus 100 mM NaCl. Final composition of the buffer is 1 mM monobasic potassium phosphate, 3 mM dibasic sodium phosphate, 253 mM sodium chloride, pH 7.4.

[0135] The column is loaded with 4 ml of FSH from DAC step in 1x PBS as described above at a flow rate: 3 ml/min collecting 1 min (3 ml) fractions.

[0136] The purity of FSH after this step is usually >95% by Coomassie and silver-stained gels.

## 5 EXAMPLE 6

### Effect Of Preservatives On Physical Stability Of FSH

[0137] Since preservatives tend to denature or destabilize protein or induce aggregation (Brange, J. and Langkjaer, L., Acta Pharm. Nord, 4, 149-158, 1992; Maa YF and Hsu C, International Journal of Pharmaceutics, 140, 155-168, 1996), the physical stability of uFSH (uFSH - Vitro Diagnostics - Human Urofollitropin) in the presence of different preservatives was examined using the dynamic light scattering technique. All measurements were obtained with a system consisting of a Lexel 95 Argon Laser (488 nm), a Brookhaven Instruments model BI-200SM goniometer, and a BI9000AT autocorrelator. Data parameters consist of: initial photon counts adjusted to 100,000 counts/sec, 30 second duration, 31 dust cutoff value, and a 90° scattering angle.

[0138] Preservatives were added to a 1.5 ml solution of 1 mg/ml urinary follicle stimulin hormone (uFSH - Vitro Diagnostics - Human Urofollitropin) which had been dialyzed against 1 x PBS overnight at pH 7.4. The concentration of preservative was selected to be the concentration generally known to provide adequate anti-microbial activity. In a laminar flow hood, this sample was filtered through a 0.1 µm Anotop-Plus filter (10mm) into a 12 mm DLS test tube. The sample was placed in the DLS holder which had been equilibrated at 37°C. The auto-correlation function was determined every 15 minutes for 24 hours and analyzed to yield the hydrodynamic parameter. This measurement demonstrated that more than 99% of the protein molecules had an average diameter of about 5.7 nm. A small population (<1%) had an average diameter of about 200 nm. The presence of the preservatives did not change the size distribution of the molecule appreciably after 24 hour at 37°C. The representative data at 24 hr. was then analyzed using NNLS (Non Negative Linear Squares) program as shown in Table I. More than 99% of the molecules were in particles with an average diameter of about 5.7 nm. Using an empirical equation relating crystallographically determined hydrodynamic radius with molecular weight (Squire, P.G. and Himmel, M.E., in Arch. Bioch. Biophys., 196, pp. 165-177, 1979) this average DLS particle size corresponds to about 36,000 daltons, which is consistent with the molecular weight of the uFSH heterodimer. The remaining small population of particles (<1%) had an average diameter of about 200 nm. These data show, in Table VI that the preservatives studied did not significantly aggregate uFSH under the conditions tested.

Table VI.

Size distribution of uFSH in formulations containing different preservatives.			
Preservative	Preservative Concentration (mg/ml)	Small Particles (~ 5.7 nm)	Large Particles (~ 200 nm)
None	0	>99 %	< 1 %
<i>m</i> -cresol	3.5	>99 %	< 1 %
Phenol	3.5	>99 %	< 1 %
Benzyl alcohol	10	>99 %	< 1 %
Methylparaben	1	>99 %	< 1 %
Chlorocresol	2	>99 %	< 1 %

## EXAMPLE 7

### Thermal Denaturation Studies on uFSH Formulations

[0139] The thermal unfolding transition for uFSH (uFSH - Vitro Diagnostics - Human Urofollitropin) as a function of solvent conditions was monitored by differential scanning calorimetry (DSC). Experiments were carried out on a VP-DSC MicroCalorimeter (MicroCal inc., Northampton, MA; Plotnik, V.V., et. al., Anal. Biochem., 250:237-244, 1997) using VPViewer software for data acquisition and Origin DSC software for data analysis. The matched sample cell and reference cell were lollipop-shaped, fabricated from tantalum, with a working volume of 0.5 ml. Approximately 1 mg/ml uFSH samples were dialyzed against appropriate buffer overnight and concentration of the protein in the sample was determined by UV spectroscopy. The proteins were then diluted to 0.4 - 0.5 mg/ml for the DSC experiments. The

dialysate was used as reference solution. Sample and reference solutions were degassed for 5 minutes before loading into the cells with a 2.5 ml needle through a filling funnel. Pressure was kept at about 30 psi with the pressure cap. For all measurements in this study, the instrument was run overnight with buffers in both the reference cell and sample cell to establish thermal history prior to sample runs. The data was analyzed with Origin DSC software using a two-state model (Sturtevant, J.M., Annu. Rev. Phys. Chem., 38:463-488, 1987). The midpoint of the transition temperature ( $T_m$ ) at different solution conditions is summarized in Table VII. The protein undergoes a very cooperative transition with a  $T_m$  of 77.3 °C in PBS buffer at pH 7.4. Thermal denaturation is irreversible on the time scale of the measurements as shown by the absence of transition in the second scan immediately following the first scan. However, the dissociated subunits stay as monomers in solution and these monomers can then reassociate to form biologically active dimer in the course of the days (data not shown). The effects of the addition of preservatives, *m*-cresol, phenol, benzyl alcohol, methylparaben, and chlorocresol at the concentrations specified below shows only marginal effect on the  $T_m$  as demonstrated in Tables VII.

Table VII.

Effect of pH, salt, and preservatives on $T_m$ of uFSH in solution as monitored by DSC.	
Solution conditions	$T_m$ (°C)
PB (9.5 mM phosphate)	
pH 5.7	72.4
pH 6.6	74.3
pH 7.6	74.8
pH 8.6	75.3
100 mM NaCl, pH 7.6	76.1
PHS at pH 7.4	77.3
3.5 mg/ml <i>m</i> -Cresol	75.3
3.5 mg/ml Phenol	75.0
10 mg/ml Benzyl Alcohol	73.5
1 mg/ml Methyl Paraben	76.1
2 mg/ml Chlorocresol	74.6

## Example 8

## Stability of uFSH as a Function of pH

**[0140]** Stability of uFSH (uFSH - Vitro Diagnostics - Human Urofollitropin) in PBS was determined for various pHs. Percent of heterodimer as a function of pH was monitored by size exclusion chromatography (SEC) in Table VIII.

Table VIII.

Percent of heterodimer as a function of pH as monitored by SEC.	
pH	Percent Dimer
7.0	95.0
6.5	95.0
6.0	95.0
5.5	95.0
5.0	95.0
4.5	95.0
4.0	95.0

Table VIII. (continued)

Percent of heterodimer as a function of pH as monitored by SEC.	
pH	Percent Dimer
3.5	95.0
3.25	87.7
3.0	62.3
2.5	17.6
2.0	5.0

## EXAMPLE 9

## Stability of uFSH Preserved and Non-Preserved Formulations

**[0141]** A solution of uFSH (uFSH - Vitro Diagnostics - Human Urofollitropin) was prepared in PBS (Dulbecco's, GIBCO) and further diluted with PBS to a concentration of about 50 µg/ml. The protein concentration was determined on a Hewlett Packard Model 8452A Diode Array Spectrophotometer.

**[0142]** A portion of the uFSH solution was added to a beaker containing a pre-weighed sample of *m*-cresol to give a final *m*-cresol concentration of about 3.16 mg/ml. 1-ml aliquots of the preserved and non-preserved solution were placed in plastic centrifuge tubes and incubated up to 238 days at about 22°C, 37°C and 45°C. At various times, aliquots were injected onto a Superdex-75 HR 10/30 column (Pharmacia) equilibrated and run at ambient temperature at 0.5 ml/min. in PBS. The eluant was monitored at 214 nm. The percentage of the heterodimer was calculated from the ratio of the area of dimer peak divided by the total area of dimer peak and monomer peaks, as shown in Table III. After 64 days, about 79% of the uFSH molecules in solution with *m*-cresol remain as intact dimer at 37°C and more than 52 % stay as dimer at 45°C. Surprisingly, after 63 days at room temperature, there is minimal dissociation of uFSH heterodimer in both the non-preserved and preserved solutions. It is remarkable that at 23°C, 37°C, and 45°C there is generally relatively low dissociation of uFSH heterodimer in both non-preserved and preserved solutions at days 4, 8, 16, 21, 28, 29, 43, 63, 64, 126, 127, 237, and 238.



Table IX. Percent Dimer in Preserved and Non-preserved uFSH solutions.

Days	PBS			PBS + <i>m</i> -cresol		
	23°C	37°C	45°C	23°C	37°C	45°C
0	96.0	96.0	96.0	93.8	93.8	93.8
4	-	93.0	91.7	-	91.8	85.6
8	94.3	92.7	89.5	93.6	88.9	80.0
16	94.1	91.1	84.3	92.5	87.0	73.8
21	94.0	91.0	83.1	-	-	-
22	-	-	-	92.3	84.7	68.8
28	92.8	89.8	-	92.7	83.7	-
29	-	-	79.9	-	-	65.3
43	93.3	89.5	75.9	91.6	81.5	59.2
63	92.9	-	-	92.6	-	-
64	-	88.2	68.6	-	78.9	52.5
126	91.5	85.5	-	89.6	71.2	-
127	-	-	58.2	-	-	43.8
237	92.3	83.1	-	91	62.0	-
238	-	-	57.9	-	-	39.2

## EXAMPLE 10

## Bioactivity Measurements of uFSH Samples

[0143] HEK 293 cells stably transfected with a cAMP sensitive b-lactamase (BLAM) reporter vector (Zlokarnik, et al., 1998, *Science* 279:84-88) were transfected with a human FSH receptor expression vector encoding a hygromycin selectable marker and incubated in hygromycin for 3 weeks. The surviving cells were treated with 10 µg/ml of FSH (Sigma) for 5 hours and the population of FSH activated cells showing the highest intensity of blue fluorescence were identified and isolated by FACS. This polyclonal population was expanded, treated with 10 µg/ml of FSH for 5 hours, and FACS sorted into single cell clones. Two clonal cell lines were analyzed by the BLAM microtiter plate assay and showed a 6 to 8 fold increase in blue/green ratio. The FSH-R cell line with the greatest fold increase in BLAM expression was chosen as the cell line to be used in the subsequent FSH assays.

[0144] The FSH receptor cell line harboring the cAMP sensitive BLAM reporter was seeded in 100 µl Growth Medium (DMEM catalog number 11965-092, 10% FBS, 500 µg/ml Gentamicin) at 20,000 cells/well in a poly-D-lysine coated, 96-well black wall tissue culture plate, and incubated overnight at 37°C under 5% CO<sub>2</sub>. The Growth Medium was replaced by 100 µl Assay Medium (DMEM catalog number 11965-092, 0.5% FBS, 500 µg/ml Gentamicin) and the plate was incubated overnight at 37°C and 5% CO<sub>2</sub>. The Assay Medium was removed and 100 µl of Assay Medium containing the indicated concentration of the FSH was then added to each well and the plate was incubated for 5 hours at 37°C under 5% CO<sub>2</sub>. 20 µl of the BLAM substrate loading, composed of 6 µl of 1 mM CCF2-AM in DMSO, 6 µl Pluronic Acid (100 µg/ml in 0.1% acetic acid DMSO) into 1 ml 2% PEG-400 and 10% ESS (Aurora Biosciences) was then added into each well. After 1 hour of incubation at room temperature, the ratio of blue (395 nm excitation/460 nm emission) to green (395 nm excitation/530 nm emission) fluorescence intensities was determined with a Cytofluor (Perseptive Biosystems, Series 4000 multi-well plate reader). The fold increase in blue/green ratio resulting from the presence of FSH was calculated by dividing each ratio by the blue/green ratio of the control sample.

[0145] A solution of urinary FSH (uFSH - Vitro Diagnostics - Human Urofollitropin) at 50 µg/ml in PBS (Sample A) was prepared as in Example 9. A portion of this solution was heated at 90°C for 10 minutes to dissociate more than 99% of the heterodimer into the two monomers (as shown by SEC analysis) and was used in this assay (Sample B) as a negative control. Another portion of the FSH solution was modified to include about 3.15 mg/ml of *m*-cresol (Sample C). The bioactivity of these three test samples was evaluated in the FSH-R 293-Cre-BLAM assay on two separate plates. The average of triplicate analyses on each plate is shown in Table X. This data shows the assay was performing

very reproducibly. It also showed that the dissociated heterodimer lost bioactivity (Sample B) and that the FSH in the formulation containing the m-cresol (Sample C) retained full bioactivity.

Table X.

Blue/Green fold increase in the FSH receptor bioactivity assay.						
Test Sample concentration (nM)	Sample A Plate 1	Sample A Plate 2	Sample B Plate 1	Sample B Plate 2	Sample C Plate 1	Sample C Plate 2
0 (control)	1.00	1.00	1.00	1.00	1.00	1.00
0.003	1.68	1.67	1.07	1.06	1.22	1.22
0.01	1.91	1.86	1.06	1.06	1.58	1.62
0.03	3.00	2.89	1.06	1.06	2.72	3.21
0.1	4.00	4.09	1.18	1.18	3.86	3.85
0.3	4.53	4.47	1.24	1.24	4.61	4.63
1	4.88	4.71	1.68	1.69	4.81	4.77
3	4.60	4.64	2.61	2.58	4.83	4.92

## EXAMPLE 11

## Bioactivity Measurements of Preserved and Non-Preserved uFSH Samples

**[0146]** Bioactivity measurements of urinary FSH(uFSH - Vitro Diagnostics - Human Urofollitropin) samples were determined in the in vitro bioactivity assay as described in Example 10. In vitro assay results of two samples uFSH in PBS at 23 °C, PBS with and without m-cresol at 23 °C after 11 month compared to a control in table XI. The data indicates that preserved and non-preserved uFSH samples maintain full biological activity after 11 months in this assay.

Table XI:

In vitro bioactivity of preserved and non-preserved uFSH solution after 11 month at 23 °C.	
Sample	Relative potency
uFSH control (50 µg/ml, PBS, fresh)	1.0
uFSH (50 µg/ml, PBS, 23 °C for 11 months)	1.27
uFSH (50 µg/ml, PBS, 3.15 mg/ml m-cresol, 23°C for 11 months)	0.86

## EXAMPLE 12

## Cartridge Compatibility of Preserved and Non-preserved rFSH Variant

**[0147]** To test the compatibility of formulated rFSH variant (alpha subunit SEQ ID NO:5; beta subunit SEQ ID NO: 11) solutions with cartridges, 4-ml solutions of the samples as listed in Table 12 were prepared from stock solutions listed below:

1.85 mg/ml rFSH variant in PBS  
 1.73 mg/ml rFSH variant in PB  
 20 mg/ml m-cresol in PBS  
 20 mg/ml m-cresol in PB  
 20% glycerol in PB  
 1 x PBS

**[0148]** After mixing, 1.7 ml of each solution was pipetted into individual cartridges with minimal head-space allowed. Two cartridges were filled for each sample. Caps were sealed on the cartridges. The cartridges were then incubated at 30°C for 20 days. After filling, the remainder of the samples were incubated at 40°C to serve as control samples.

**[0149]** In vitro activity of these samples was measured after 20 days of incubation at 30°C, using the method de-

scribed in Example 11. The activity of these samples was compared to corresponding control samples at zero time point. As shown in Table XII below, rFSH sample at 50 µg/ml in PBS and sample at 200 µg/ml in PBS and 3.15 mg/ml m-cresol are stable in cartridges (cartridge 1 and cartridge 4). These samples remain fully active after 20 days of incubation at 30°C. However, samples in phosphate buffer without NaCl were less potent under these conditions. The activity of rFSH samples in phosphate buffer and 1.6 % glycerol decreases in comparison to that of control (cartridge 2 and cartridge 3)

Table XII.

In vitro activity of preserved and non-preserved samples incubated at 30 °C for 20 days in cartridges.		
Sample	Sample Conditions	Relative Potency
Cartridge 1	200 µg/ml rFSH variant, 3.15 mg/ml m-cresol, PBS, pH 7.4	1.06
Cartridge 2	200 µg/ml rFSH variant, 1.6% glycerol, PB, pH 7.4	0.81
Cartridge 3	50 µg/ml rFSH variant, 1.6% glycerol, 3.15 mg/ml m-cresol, PB, pH 7.4	0.60
Cartridge 4	50 pg/ml rFSH variant, PBS, pH 7.4	1.10

As demonstrated by these examples, following the methods and techniques described one can generate surprisingly stable composition and formulations of FSH or a FSH variant. These compositions and formulations result in the development of the presently claimed articles of manufacture. Since about 1970, the courts have held that printed information in an article of manufacture does not remove the article from the realm of patentability so long as the item and the invention as a whole satisfy the other requirements of the statute, such as novelty and nonobviousness. Since the FSH or a FSH variant products taught in the prior art expressly teach that such solutions are suitable only for immediate use and after use the contents must be disposed of, rather than the presently claimed product suitable for use 24 hours or greater, the article of manufacture embodies new and non-obvious invention that is distinct and different from the prior art.

**[0150]** The principles, preferred embodiments, and modes of operation of the present invention have been described in the foregoing specification. The invention intended to be protected herein, however, is not to be construed as limited to the particular forms disclosed, since they are to be regarded as illustrative rather than restrictive. Variations and changes may be made by those skilled in the art without departing from the spirit of the invention.

## EXAMPLE 9

## Formulation Stability of Preserved and Non-Preserved FSH Variant Samples

**[0151]** A stock solution of recombinant FSH variant (alpha subunit is SEQ ID:5; beta subunit SEQ ID:11) at about 1 mg/ml in phosphate buffered saline (PBS, Dulbecco's, GIBCO) was diluted to 50 µg/ml or 20 µg/ml with either PBS or PBS containing m-cresol to give a final m-cresol concentration of 3.15 mg/ml. Similarly, another set of samples was made using 10 mg/ml benzyl alcohol as preservative. 1-ml aliquots of the preserved and non-preserved solution were incubated at 4, 22, 37 °C for up to three months in plastic eppendorf tubes. At various times, aliquots were injected onto a Superdex 75 gel filtration column (Pharmacia) equilibrated with PBS and run at ambient temperature with a flow rate of 0.07 ml/min and the run time of 35 minutes. Detection was monitored by UV absorbance at 214 nm over time. Peak areas were integrated, and the percentage of the heterodimer was calculated as a ratio of area of heterodimer peak over the total area of the dimer and monomer peaks.

As shown in Table IX, there is minimal dissociation of the heterodimer under various solution conditions after three month of incubation at room temperature or below. Greater than 50% heterodimer remains intact after three month incubation at 37°C. The stability of heterodimer is higher with more concentrated solution.

Table IX.

Heterodimer stability of rFSH variant monitored by SE-HPLC.						
Sample	% Dimer					
	1 month			3 months		
	4°C	22°C	37°C	4°C	22°C	37°C
20 µg/ml in PBS	100	100	88.9	100	100	77.3

Table IX. (continued)

Heterodimer stability of rFSH variant monitored by SE-HPLC.						
Sample	% Dimer					
	1 month			3 months		
	4°C	22°C	37°C	4°C	22°C	37°C
20 µg/ml in PBS 3.15 mg/ml m-cresol	100	100	86.2	100	100	64.6
20 µg/ml in PBS 10 mg/ml benzyl alcohol	100	100	89.1	100	100	57.1
50 µg/ml in PBS	100	100	100	100	100	81.1
50 µg/ml in PBS 3.15 mg/ml m-cresol	100	100	90	100	100	75.7
50 µg/ml in PBS 10 mg/ml benzyl alcohol	100	100	87	100	100	61.0

Annex to the description

[0152]

SEQUENCE LISTING

5

<110> Eli Lilly and Company

10

<120> FSH AND FSH VARIANT FORMULATIONS, PRODUCTS AND METHODS

<130> X12383M Sequence Listing

15

<140>

<141>

20

<150> 60/093906

<151> 1998-07-23

25

<150> 60/094611

<151> 1998-07-30

30

<150> 60/094767

<151> 1998-07-31

35

<150> 60/098711

<151> 1998-09-01

40

<150> 60/100696

<151> 1998-09-17

45

<160> 20

50

<170> PatentIn Ver. 2.0

55

<210> 1

EP 0 974 359 A2

<211> 96

<212> PRT

5

<213> mammalian

<400> 1

10

Phe Pro Asp Gly Glu Phe Thr Met Gln Gly Cys Pro Glu Cys Lys Leu

1 5 10 15

15

Lys Glu Asn Lys Tyr Phe Ser Lys Pro Asp Ala Pro Ile Tyr Gln Cys

20 25 30

20

Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Ala Arg Ser Lys

35 40 45

25

Lys Thr Met Leu Val Pro Lys Asn Ile Thr Ser Glu Ala Thr Cys Cys

50 55 60

30

Val Ala Lys Ala Phe Thr Lys Ala Thr Val Met Gly Asn Val Arg Val

65 70 75 80

35

Glu Asn His Thr Glu Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser

85 90 95

40

45

50

<210> 2

<211> 111

55

<212> PRT

EP 0 974 359 A2

<213> mammalian

5

<400> 2

Arg Ser Cys Glu Leu Thr Asn Ile Thr Ile Thr Val Glu Lys Glu Glu

1 5 10 15

10

Cys Gly Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys

20 25 30

15

Tyr Thr Arg Asp Leu Val Tyr Arg Asp Pro Ala Arg Pro Asn Ile Gln

35 40 45

20

Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Lys Val Pro

50 55 60

25

Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr

65 70 75 80

30

Glu Cys His Cys Ser Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val

85 90 95

35

Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Arg Glu Ile Lys Glu

100 105 110

40

45

<210> 3

<211> 96

50

<212> PRT

<213> mammalian

55

<400> 3

EP 0 974 359 A2

Phe Pro Asp Gly Glu Phe Thr Thr Gln Asp Cys Pro Glu Cys Lys Leu  
 1 5 10 15  
 5

Arg Glu Asn Lys Tyr Phe Phe Lys Leu Gly Val Pro Ile Tyr Gln Cys  
 20 25 30  
 10

Lys Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Ala Arg Ser Arg  
 35 40 45  
 15

Lys Thr Met Leu Val Pro Lys Asn Ile Thr Ser Glu Ser Thr Cys Cys  
 50 55 60  
 20

Val Ala Lys Ala Phe Ile Arg Val Thr Val Met Gly Asn Ile Lys Leu  
 65 70 75 80  
 25

Glu Asn His Thr Gln Cys Tyr Cys Ser Thr Cys Tyr His His Lys Ile  
 85 90 95  
 30

35

40

<210> 4

<211> 111  
 45

<212> PRT

<213> mammalian

50

<400> 4

Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Val Glu Lys Glu Gly  
 1 5 10 15  
 55



EP 0 974 359 A2

Cys Gly Phe Cys Ile Thr Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys  
 20 25 30  
 5  
 Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Asn Ile Gln  
 35 40 45  
 10  
 Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Lys Val Pro  
 50 55 60  
 15  
 Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr  
 65 70 75 80  
 20  
 Ala Cys His Cys Gly Lys Cys Asn Ser Asp Ser Thr Asp Cys Thr Val  
 85 90 95  
 25  
 Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Asp Met Lys Glu  
 100 105 110  
 30  
 35  
 <210> 5  
 <211> 92  
 40  
 <212> PRT  
 <213> Homo sapiens  
 45  
 <400> 5  
 Ala Pro Asp Val Gln Asp Cys Pro Glu Cys Thr Leu Gln Glu Asn Pro  
 1 5 10 15  
 50  
 Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys  
 20 25 30  
 55

EP 0 974 359 A2

Phe Ser Arg Ala Tyr Pro Thr Pro Leu Arg Ser Lys Lys Thr Met Leu  
 5 35 40 45  
 Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser  
 10 50 55 60  
 Tyr Asn Arg Val Thr Val Met Gly Gly Phe Lys Val Glu Asn His Thr  
 15 65 70 75 80  
 Ala Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser  
 20 85 90  
 25  
 <210> 6  
 <211> 111  
 30 <212> PRT  
 <213> Homo sapiens  
 35  
 <400> 6  
 Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu  
 40 1 5 10 15  
 Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys  
 45 20 25 30  
 Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln  
 50 35 40 45  
 Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro  
 55

EP 0 974 359 A2

	50	55	60
5	Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr		
	65	70	75 80
10	Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val		
	85	90	95
15	Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu Met Lys Glu		
	100	105	110
20			
	<210> 7		
25	<211> 96		
	<212> PRT		
	<213> mammalian		
30			
	<400> 7		
	Phe Pro Asp Gly Glu Phe Thr Met Gln Gly Cys Pro Glu Cys Lys Leu		
35	1	5	10 15
	Lys Glu Asn Lys Tyr Phe Ser Lys Leu Gly Ala Pro Ile Tyr Gln Cys		
40	20	25	30
	Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Ala Arg Ser Lys		
45	35	40	45
	Lys Thr Met Leu Val Pro Lys Asn Ile Thr Ser Glu Ala Thr Cys Cys		
50	50	55	60
	Val Ala Lys Ala Phe Thr Lys Ala Thr Val Met Gly Asn Ala Arg Val		
55			

EP 0 974 359 A2

65 70 75 80

5

Glu Asn His Thr Glu Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser

85 90 95

10

15

20 <210> 8  
<211> 111  
<212> PRT

25 <213> mammalian

30 <400> 8

Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Thr Val Glu Lys Glu Glu

1 5 10 15

35

Cys Asn Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys

20 25 30

40

Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Asn Ile Gln

35 40 45

45

Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Lys Val Pro

50 50 55 60

55

Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr

65 70 75 80

EP 0 974 359 A2

Glu Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val  
 85 90 95  
 5  
 Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Ser Glu Met Lys Glu  
 100 105 110  
 10  
 <210> 9  
 <211> 96  
 <212> PRT  
 20  
 <213> mammalian  
 <400> 9  
 25  
 Phe Pro Asp Gly Glu Phe Thr Met Gln Gly Cys Pro Glu Cys Lys Leu  
 1 5 10 15  
 30  
 Lys Glu Asn Lys Tyr Phe Ser Lys Pro Asp Ala Pro Ile Tyr Gln Cys  
 20 25 30  
 35  
 Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Ala Arg Ser Lys  
 35 40 45  
 40  
 Lys Thr Met Leu Val Pro Lys Asn Ile Thr Ser Glu Ala Thr Cys Cys  
 50 55 60  
 45  
 Val Ala Lys Ala Phe Thr Lys Ala Thr Val Met Gly Asn Val Arg Val  
 65 70 75 80  
 50  
 Glu Asn His Thr Glu Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser  
 85 90 95  
 55

<210> 10  
 5 <211> 111  
 <212> PRT  
 10 <213> mammalian  
  
 <400> 10  
 15 Arg Ser Cys Glu Leu Thr Asn Ile Thr Ile Thr Val Glu Lys Glu Glu  
     1                    5                    10                    15  
  
 20 Cys Ser Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys  
                     20                    25                    30  
  
 25 Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Asn Ile Gln  
                     35                    40                    45  
  
 30 Lys Ala Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Lys Val Pro  
                     50                    55                    60  
  
 35 Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr  
 40 65                    70                    75                    80  
  
 Glu Cys His Cys Gly Lys Cys Asp Arg Asp Ser Thr Asp Cys Thr Val  
 45                    85                    90                    95  
  
 Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Ser Asp Ile Arg Glu  
 50                    100                    105                    110  
  
 55

EP 0 974 359 A2

<210> 11

5 <211> 108

<212> PRT

10 <213> Homo sapiens

<400> 11

15 Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu

1 5 10 15

20 Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys

20 25 30

25 Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln

35 40 45

30 Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro

50 55 60

35 Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr

65 70 75 80

40 Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val

85 90 95

45 Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu

50 100 105

55 <210> 12

EP 0 974 359 A2

5 <211> 109  
 <212> PRT  
 <213> Homo sapiens  
  
 10 <400> 12  
 Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu  
 1 5 10 15  
 15 Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys  
 20 20 25 30  
 Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln  
 25 35 40 45  
 Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro  
 30 50 55 60  
 Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr  
 35 65 70 75 80  
 Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val  
 40 85 90 95  
 Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu Met  
 45 100 105  
  
 50 <210> 13  
 <211> 110  
 55 <212> PRT



<213> Homo sapiens

5

<400> 13

Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu

10

1 5 10 15

Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys

15

20 25 30

Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln

20

35 40 45

Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro

25

50 55 60

Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr

30

65 70 75 80

Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val

35

85 90 95

Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu Met Lys

40

100 105 110

45

<210> 14

<211> 276

50

<212> DNA

<213> Homo sapiens

55

<400> 14

EP 0 974 359 A2

gctcctgatg tgcaggattg cccagaatgc acgctacagg aaaacccatt cttctcccag 60  
 5 cccgggtgcc caatacttca gtgcatgggc tgctgcttct ctagagcata tccactcca 120  
 ctaagggtcca agaagacgat gttggtccaa aagaacgtca cctcagagtc cacttgctgt 180  
 gtagctaaat catataacag ggtcacagta atgggggggt tcaaagtga gaaccacacg 240  
 10 gcgtgccact gcagtacttg ttattatcac aaatct 276

<210> 15

15

<211> 324

<212> DNA

<213> Homo sapiens

20

<400> 15

25

aatagctgtg agctgaccaa catcaccatt gcaatagaga aagaagaatg tcgtttctgc 60  
 ataagcatca acaccacttg gtgtgctggc tactgctaca ccagggatct ggtgtataag 120  
 gaccagcca ggccaaaat ccagaaaaca tgtaccttca aggaactggg atatgaaaca 180  
 30 gtgagagtgc ccggctgtgc tcaccatgca gattccttgt atacataccc agtggccacc 240  
 cagtgtcact gtggcaagtg tgacagcgac agcactgatt gtactgtgcg aggcctgggg 300  
 cccagctact gtccttttgg tgaa 324

35

<210> 16

<211> 327

40

<212> DNA

<213> Homo sapiens

45

<400> 16

aatagctgtg agctgaccaa catcaccatt gcaatagaga aagaagaatg tcgtttctgc 60  
 50 ataagcatca acaccacttg gtgtgctggc tactgctaca ccagggatct ggtgtataag 120  
 gaccagcca ggccaaaat ccagaaaaca tgtaccttca aggaactggg atatgaaaca 180  
 gtgagagtgc ccggctgtgc tcaccatgca gattccttgt atacataccc agtggccacc 240  
 55 cagtgtcact gtggcaagtg tgacagcgac agcactgatt gtactgtgcg aggcctgggg 300

EP 0 974 359 A2

cccagctact gctcctttgg tgaaatg

327

5

<210> 17

<211> 330

10

<212> DNA

<213> Homo sapiens

15

<400> 17

aatagctgtg agctgaccaa catcaccatt gcaatagaga aagaagaatg tcgtttctgc 60  
 ataagcatca acaccacttg gtgtgctggc tactgctaca ccagggatct ggtgtataag 120  
 gaccagcca ggccaaaat ccagaaaaca tgtaccttca aggaactggt atatgaaaca 180  
 gtgagagtgc ccggctgtgc tcacatgca gattccttgt atacataccc agtggccacc 240  
 cagtgtcact gtggcaagtg tgacagcgac agcactgatt gtactgtgcg aggcctgggg 300  
 cccagctact gctcctttgg tgaaatgaaa 330

25

<210> 18

30

<211> 333

<212> DNA

<213> Homo sapiens

35

<400> 18

aatagctgtg agctgaccaa catcaccatt gcaatagaga aagaagaatg tcgtttctgc 60  
 ataagcatca acaccacttg gtgtgctggc tactgctaca ccagggatct ggtgtataag 120  
 gaccagcca ggccaaaat ccagaaaaca tgtaccttca aggaactggt atatgaaaca 180  
 gtgagagtgc ccggctgtgc tcacatgca gattccttgt atacataccc agtggccacc 240  
 cagtgtcact gtggcaagtg tgacagcgac agcactgatt gtactgtgcg aggcctgggg 300  
 cccagctact gctcctttgg tgaaatgaaa gaa 333

50

<210> 19

<211> 276

55

<212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: Modified to  
facilitate cloning.

10

<400> 19

15

gctcctgatg tgcaggattg cccagaatgc acgctacagg aaaaccatt cttctcccag 60  
ccgggtgccc caatacttca gtgcatgggc tgctgcttct caagagcata tccactcca 120  
ctaagggtcca agaagacgat gttggtccaa aagaacgtca cctcagagtc cacttgctgt 180  
gtagctaaat catataacag ggtcacagta atgggggggt tcaaagtga gaaccacacg 240  
gcgtgccact gcagtacttg ttattatcac aaatct 276

20

25

<210> 20

<211> 324

<212> DNA

30

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence: Modified to  
facilitate cloning.

40

<400> 20

aacagctgtg agctcaccaa catcaccatt gcaatagaga aagaagaatg tcgtttctgc 60  
atatcgatca acaccacttg gtgtgctggc tactgctaca ccagggatct ggtgtataag 120  
gacccggccc gtcccaaat ccagaaaaca tgtaccttca aggaactggg atatgaaaca 180  
gtacgcgtgc ccggctgtgc tcaccatgca gattccttgt atacataccc agtggccacc 240  
cagtgtcact gtggcaagtg tgacagcgac agcactgatt gtactgtgcg aggcctgggg 300  
cccagctact gtcctttgg tgaa 324

45

50

55

## Claims

1. A formulation comprising FSH or a FSH variant, containing an alpha and beta subunit, and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent.
2. A formulation of Claim 1, wherein the preservative is phenol, m-cresol, chlorocresol, or a mixture thereof.
3. A formulation of Claim 2, wherein the concentration of FSH or a FSH variant is about 1.0 µg/ml to about 50 mg/ml.
4. A formulation of Claim 3, further comprising an isotonicity agent.
5. A formulation of Claim 4, further comprising a physiologically acceptable buffer.
6. A formulation comprising FSH or a FSH variant lyophilized in a first vial, and a second vial containing a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent.
7. A formulation of Claim 1, wherein said FSH or a FSH variant and preservative are in solution.
8. A formulation of Claim 1, wherein said FSH or a FSH variant is at least one compound selected from the group consisting of:

(a) :  $\alpha$ -subunit: (SEQ ID NO:1)

FPDGEFTMQGCPECKLKENKYFSKPDAPYIYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:2)

RSCELTNITITVEKEECGFCISINTTWCAGYCYTRDLVYRDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCSKCDSDSTDCTVRGLGPSYCSFREIKE

(b) :  $\alpha$ -subunit: (SEQ ID NO:3)

FPDGEFTTQDCPECKLRENKYFFKLGVPYIYQCKGCCFSRAYPTPARSRKTMLVPKN  
ITSESTCCVAKAFIRVTVMGNIKLENHTQCYCSTCYHHKI

$\beta$ -subunit: (SEQ ID NO:4)

NSCELTNITIAVEKEGCGFCITINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATACHCGKCNSDSTDCTVRGLGPSYCSFGDMKE

(c): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:6)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMKE

(d): $\alpha$ -subunit:(SEQ ID NO:7)

FPDGEFTMQGCPECKLKENKYFSKLGAPIYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNARVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:8)

NSCELTNITITVEKEECNFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCGKCDSDSTDCTVRGLGPSYCSFSEMKE

(e): $\alpha$ -subunit:(SEQ ID NO:9)

FPDGEFTMQGCPECKLKENKYFSKPDAPYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:10)

RSCELTNITITVEKEECNFCISINTTWCAGYCYTRDLVYKDPARPNIQACTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCGKCDRDSTDCTVRGLGPSYCSFSDIRE

(f): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:11)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGE

(g) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:12)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEM

(h) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:13)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMK

9. A process for preparing a preserved solution formulation of FSH or a FSH variant, containing an alpha and beta subunit, which comprises admixing said FSH or a FSH variant and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof, in an aqueous diluent.
10. An article of manufacture for human pharmaceutical use, comprising packaging material and a vial comprising a solution of FSH or a FSH variant, containing an alpha and beta subunit, and a preservative solution, wherein said packaging material comprises a label which indicates that said solution may be held over a period of 24 hours or greater.
11. The article of manufacture of Claim 10, wherein said vial is a glass container having a stopper for multi-use administration.
12. The article of manufacture of Claim 10, wherein said vial is a blister pack, capable of being punctured and used in pulmonary administration.
13. The article of manufacture of Claim 10, wherein said vial is a pen-injector device.
14. An article of manufacture, comprising packaging material, a first vial comprising lyophilized FSH or a FSH variant, containing an alpha and beta subunit, and a second vial comprising a preservative solution, wherein said packaging material comprises a label which instructs a patient to reconstitute the said lyophilized FSH or a FSH variant in the preservative solution for use over a period of 24 hours or greater.
15. The article of manufacture of Claim 14, wherein said first vial and said second vial are embodied in a pen-injector device.
16. The use of at least one alpha or beta polypeptide of a FSH or a FSH variant in the preparation of a preserved formulation adapted for administration over a period of 24 hours or greater.
17. A stable formulation comprising at least one FSH or a FSH variant, containing an alpha and beta subunit, and phosphate buffer containing saline or a salt, wherein said FSH or a FSH variant comprises at least 90% FSH or a FSH variant dimers after 60 days at 23°C.

18. A formulation of Claim 17, wherein the concentration of said FSH or a FSH variant is about 1.0 µg/ml to about 50 mg/ml.

19. A formulation of Claim 17, further comprising an isotonicity agent.

20. A formulation of Claim 17, wherein said buffer is at least one physiologically acceptable buffer.

21. A formulation comprising a first vial containing a lyophilized FSH or a FSH variant containing an alpha and beta subunit, and a second vial containing phosphate buffer containing saline or a salt.

22. A formulation of Claim 17, wherein said at least one said FSH or a FSH variant and at least one said buffer are in solution.

23. A formulation of Claim 17, wherein said FSH or a FSH variant is at least one compound selected from the group consisting of:

(a) :  $\alpha$ -subunit: (SEQ ID NO:1)

FPDGEFTMQGCPECKLKENKYFSKPDAPIYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:2)

RSCELTNITITVEKEECGFCISINTTWCAGYCYTRDLVYRDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCSKCDSDSTDCTVRGLGPSYCSFREIKE

(b) :  $\alpha$ -subunit: (SEQ ID NO:3)

FPDGEFTTQDCPECKLRENKYFFKLGVPYIYQCKGCCFSRAYPTPARSRKTMLVPKN  
ITSESTCCVAKAFIRVTVMGNIKENHTQCYCSTCYHHKI

$\beta$ -subunit: (SEQ ID NO:4)

NSCELTNITIAVEKEGCGFCITINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATACHCGKCNDSSTDCTVRGLGPSYCSFGDMKE

(c) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:6)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMKE



(d): $\alpha$ -subunit:(SEQ ID NO:7)

FPDGFTMQGCPECKLKENKYFSKLGAPIYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNARVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:8)

NSCELTNITITVEKEECNFCISINTTWCAGYCYTRDLVYKDPARPNIQKTCTFKEL  
VYETVKVPGCAHHADSLYTPVATECHCGKCDSSTDCTVRGLGPSYCSFSEMKE

(e): $\alpha$ -subunit:(SEQ ID NO:9)

FPDGFTMQGCPECKLKENKYFSKPDAPYQCMGCCFSRAYPTPARSKKTMLVPKN  
ITSEATCCVAKAFTKATVMGNVRVENHTECHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:10)

RSCELTNITITVEKEECSCFCISINTTWCAGYCYTRDLVYKDPARPNIQKACTFKE  
LVYETVKVPGCAHHADSLYTPVATECHCGKCDRDSTDCTVRGLGPSYCSFSDIR  
E

(f): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:11)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSSTDCTVRGLGPSYCSFGE

(g): $\alpha$ -subunit:(SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit:(SEQ ID NO:12)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSSTDCTVRGLGPSYCSFGEM

(h) :  $\alpha$ -subunit: (SEQ ID NO:5)

APDVQDCPECTLQENPFFSQPGAPILQCMGCCFSRAYPTPLRSKKTMLVQKNVTSE  
STCCVAKSYNRVTVMGGFKVENHTACHCSTCYHKS

$\beta$ -subunit: (SEQ ID NO:13)

NSCELTNITIAIEKEECRFCISINTTWCAGYCYTRDLVYKDPARPKIQKTCTFKEL  
VYETVRVPGCAHHADSLYTPVATQCHCGKCDSDSTDCTVRGLGPSYCSFGEMK

24. A process for preparing a stable solution formulation of FSH or a FSH variant containing an alpha and beta subunit, which comprises admixing a FSH or a FSH variant in phosphate buffer containing saline or a salt.
25. An article of manufacture for pharmaceutical use, comprising packaging material and a vial comprising a stable solution of FSH or a FSH variant containing an alpha and beta subunit in an aqueous diluent, wherein said packaging material comprises a label which indicates that such solution is suitable for use over a period of 24 hours or greater.
26. The article of manufacture of Claim 25, wherein said vial is a glass container having a stopper for multi-use administration.
27. The article of manufacture of Claim 25, wherein said vial is a blister pack, capable of being punctured and used in pulmonary administration.
28. The article of manufacture of Claim 25, wherein said vial is a pen-injector device.
29. An article of manufacture, comprising packaging material, a first vial comprising a lyophilized FSH or a FSH variant containing, an alpha and beta subunit, and a second vial comprising a stable aqueous diluent, wherein said packaging material comprises a label which instructs a patient to reconstitute said FSH or a FSH variant in the aqueous diluent to form a solution that is suitable for use over a period of 24 hours or greater.
30. The article of manufacture of Claim 29, wherein said first vial and said second vial are embodied in a pen-injector device.
31. The use of at least one polypeptide of a FSH or a FSH variant in the preparation of a non-preserved formulation adapted for administration over a period of 24 hours or greater.
32. Use of a formulation of claim 1 for treating infertility in a patient in need thereof.
33. Use of a formulation of claim 1 wherein said patient is selected from the group consisting of a human, sheep, cow, pig, horse, or rabbit.
34. Use of a preserved solution of FSH or a FSH variant, containing an alpha and beta subunit, to treat infertility in a patient in need thereof, said solution being suitable for administration over a period of 24 hours or greater.
35. Use of a stable solution of FSH or a FSH variant, containing an alpha and beta subunit, to treat infertility in a patient, which comprises administering to a patient in need thereof a solution of said FSH or a FSH variant in a phosphate buffer, containing saline or a salt, over a period of 24 hours or greater.
36. Use of a formulation of Claim 17 for treating infertility in a patient in need thereof.
37. A use of Claim 41 wherein said patient is selected from the group consisting of a human, sheep, cow, pig, horse, or rabbit.
38. Use of a stable solution of purified FSH or a FSH variant, containing an alpha and beta subunit, in a phosphate buffer containing saline or a salt suitable for administration over a period of 24 hours or greater for treating infertility

in a patient in need thereof.

39. Use of a stable solution FSH or a FSH variant, containing an alpha and beta subunit, to treat infertility in a patient in need thereof, wherein said non-preserved solution of said FSH or a FSH variant in phosphate buffer containing saline or a salt is suitable for use over a period of 24 hours or greater.

5

10

15

20

25

30

35

40

45

50

55